



# 2011 Sunflower Planting Date Study



Dr. Heather Darby, UVM Extension Agronomist  
Hannah Harwood, Erica Cummings, and Rosalie Madden  
University of Vermont Crop and Soils Technicians  
(802) 524-6501

Visit us on the web at <http://www.uvm.edu/extension/cropsoil>

**2011 Vermont Sunflower Planting Date Study**  
**Heather Darby, University of Vermont Extension**  
**heather.darby@uvm.edu**

Planting date and variety selection are practices implemented to maximize yield and often crop quality. Optimal planting dates for sunflowers have yet to be determined for the Northeast. In Vermont, farmers generally try to have their sunflower crop planted by late May to meet the long growing season required by this crop. Based on research conducted in other regions, modifying planting date and variety may also help reduce bird, insect, and disease pressure. Therefore the goal of this project was to determine the impact of planting date and variety selection on sunflower yield and pest pressure. While the data presented is only representative of one year, this information can be combined with other research to aid in making planting date decisions for sunflowers in the Northeast.

## MATERIALS AND METHODS

To evaluate the impact of planting date on sunflower yield and quality, a research trial was initiated at Borderview Farm in Alburgh, VT. Agronomic information for trial can be found in Table 1. The experimental design was a randomized complete block with split plots replicated three times. The plot size was 5'x20'. The main plots were 3 planting dates (25-May, 1-Jun, and 7-Jun). The subplots were a long and a short season sunflower variety. Croplan Genetics variety '306' had a relative maturity of 87 days and Syngenta variety '7120' a relative maturity of 95 days. The soil was a Benson rocky silt loam and plots prepared by with spring disking, harrowing, and finished with a spike-tooth harrow. A starter fertilizer with an analysis of 10-20-20 was applied at a rate of 260 lbs per acre at planting. Weeds were managed by a preplant application of Treflan® (trifluralin) at 2.5 pints per acre. For postemergence weed control plots were hand-weeded (16-Jun) and cultivated with a Brillion 4-row cultivator (27-Jun).

**Table 1. Agronomic practices for the 2011 sunflower planting date study at Borderview Farm.**

<b>Location</b>	Borderview Farm – Alburgh, VT
<b>Soil type</b>	Benson rocky silt loam
<b>Previous crop</b>	Corn Silage
<b>Tillage operations</b>	Spring disk, harrow, spike-toothed harrow
<b>Weed control</b>	Trifluralin, pre-plant, 2.5 pints/acre Hand-weeded and row cultivated
<b>Seeding rate</b>	32,000 plants/acre
<b>Row width (in.)</b>	30
<b>Planting dates</b>	25-May, 1-Jun, 7-Jun
<b>Starter fertilizer (at planting)</b>	260 lbs/acre, 10-20-20
<b>Harvest dates</b>	26-Sep and 12-Oct

In late July and early August, sunflowers began to flower, and when at least 75% of a given plot was in bloom, the date was noted. During the season, bird netting was used to discourage birds from damaging the developed sunflower seed heads. With this method, bird damage was kept to a minimum. Prior to harvest sunflower population, height and head width, as well as the incidence of lodging and bird damage was recorded. Bird damage was estimated using percent evaluations provided by North Dakota State University Extension. Incidence of white mold (*Sclerotinia sclerotiorum*) at three locations on the plant: on the sunflower head, along the stalk, and at the base.



Plots were harvested on September 26 with an Almaco SP50 plot combine with a 5' head and custom-made sunflower pans (Figure 1). The 7-June planted date treatment was harvested on 12-October. Following harvest the test weight was measured with a Berckes Test Weight Scale and a Dickey-John M20P moisture meter was used to measure harvest moisture levels. Harvested seeds were then cleaned with a Clipper fanning mill and then evaluated for insect damage. Banded sunflower moth (*Cochylis hospes*) larvae have been found in the region, and are evidenced by round exit holes in seeds, caused by larvae burrowing in to feed (Figure 2). Banded sunflower moth can cause significant yield losses.

**Figure 1. Almaco SP50 plot combine at Borderview Farm on harvest day.**

Prior to oil extraction, seed samples were dried and moisture levels quantified. Oil was extruded from a

subsample of each harvested plot using a Kern Kraft Oil Press KK40. After pressing, oil content and yields were determined.

Variations in yield and quality can occur because of variations in genetics, soil, weather, and other growing conditions. Statistical analysis makes it possible to determine whether a difference among treatments is real or whether it might have occurred due to other variations in the field. All data was analyzed using a mixed model analysis where replicates were considered random effects. At the bottom of each table a Least Significant Difference (LSD) value is presented for each variable (e.g. plant height). LSDs at the 10% level (0.10) of probability are shown. Where the difference between two treatments within a column is equal to or greater than the LSD value at the bottom of the column, you can be sure in 9 out of 10 chances that there is a real difference between the two values. Treatments that were not significantly lower in performance than the highest value in a particular column are indicated with an asterisk.

In the example at right, treatment C is significantly different from treatment A but not from treatment B. The difference between B and C is equal to 0.5, which is less than the LSD value of 2.7. This means that these treatments did not differ in yield. The difference between A and C is equal to 6.9, which is greater than the LSD value of 2.7. This means that the heights of these two treatments were significantly different from one another.

Planting date	Height
A	48.5
B	54.9*
C	55.4*
LSD (0.10)	2.7



**Figure 2. Banded sunflower moth larvae.**

## RESULTS

Using data from a weather station in close proximity to Borderview Farm in Alburgh, VT, weather data is summarized in Table 2. The 2011 growing season was wetter than normal, with very heavy precipitation in the spring and late summer. However, the months of June and July were close to average in overall rainfall, and temperatures were near normal. There were an accumulated 2,998 Growing Degree Days (GDDs) at a base temperature of 44°F; this was 238 more than the 30-year average.

**Table 2. Summarized weather data for 2011 – Alburgh, VT.**

	May	June	July	August	September	October
Average Temperature (°F) ±	58.7	67.1	74.4	70.4	63.8	51.5
Departure from Normal	2.1	1.3	3.3	1.6	5.8	4.5
Precipitation (inches) *	8.67	3.52	3.68	10.23	5.56	2.68
Departure from Normal	5.35	0.09	-0.29	6.38	2.10	0.10
Growing Degree Days (base 44° F)	454	716	942	749	591	434
Departure from Normal	63.6	62.1	103.9	-26.3	98.6	241.8


± Average temperature for August-September is taken from Burlington, VT.

\* Precipitation for May-July is taken from Burlington, VT.

Based on National Weather Service data from cooperative observation stations in South Hero. Historical averages are for 30 years of data (1971-2000).

### *Planting Date by Variety Interactions*

With the exception of oil yield, there were no interactions between sunflower planting date and variety. This suggests that the varieties performed similarly across planting dates. There was an interaction between planting date and variety for oil yield. The early season variety ‘306’ had a lower oil yield than 7120 when planted 1-June and 6-June planting dates. The longer season variety ‘7120’ has higher oil yields when planted on the earliest planting date. We would expect that the shorter season variety may out perform a later season variety as the planting date becomes delayed. Likewise we would expect a later season variety to outperform a shorter season variety at earlier planting dates. Further research needs to be conducted across a broader span of planting dates to confirm these predictions.



**Figure 7. Effect of planting date and variety on oil yield.**

### Effect of Planting Date

As expected the earliest planted sunflowers (25-May) bloomed by 27-July. Sunflowers planted on 1-June bloomed by 2-August and the latest planted sunflower by the 8-August. Essentially for every one week delay in planted there was approximately a week delay in bloom.

The 1-June planting date resulted in a significantly higher plant population compared to the other dates (Table 3). Having all been seeded at a rate of 32,000 seeds per acre, this means that the establishment rates for the three planting dates varied from 53% to 81%. Sunflowers planted on the third planting date (7-June) were the tallest (55.4 inches) and also had the widest seed head (7.5 inches).


**Table 3. Effect of planting date on agronomic characteristics of sunflowers.**

Planting date	Population plants / acre	Height in	Head width in	Bird damage %	Lodging %	White mold incidence		
						Head rot %	Stalk rot %	Base rot %
5/25/2011	20600	48.5	5.9	23.5	<b>1.7</b>	<b>18.3</b>	<b>6.7</b>	<b>3.3</b>
6/1/2011	<b>25900*</b>	54.9*	5.5	15.1	13.3	26.7	18.3	<b>3.3</b>
6/7/2011	17100	<b>55.4*</b>	<b>7.5</b>	<b>6.6</b>	<b>1.7</b>	33.3	15.0	<b>3.3</b>
LSD (0.10)	3510	2.7	0.8	7.5	4.5	NS	6.9	NS
Trial Mean	21200	53.0	6.3	15.1	5.6	26.1	13.3	3.3

\* Treatments indicated with an asterisk did not perform significantly lower than the top-performing treatment in a particular column.

NS – No significant difference was determined between treatments.

Bird damage was lowest among the sunflowers planted on 7-June (6.6%) (Table 3; Figure 3). Lodging was significantly higher at the 1-June planting date than other planting dates. The incidence of white mold in the form of stalk rot was 6.7% in the first planting date (May 25), significantly lower than the second and third planting dates, which had 18.3% and 15.0% stalk rot, respectively (Table 3). There was no significant difference among planting dates for the incidence of white mold as either head rot or base rot.



**Figure 3. Effect of planting date on incidence of lodging and bird damage. Treatments with the same letter did not differ statistically ( $p=0.10$ ; compare capital letters for lodging and lower-case letters for bird damage).**

At harvest, seed moisture content was highest for the 7-June planting date (Table 4). The average moisture level was 21.0% for this third planting date, significantly higher than earlier planting dates. Seed yield was also highest for the third planting date of sunflowers (1540 lbs per acre), though not significantly higher than sunflowers planted on earlier dates. Test weight, a measure of seed density and also the amount of debris or trash in the yield, was highest in the third planting

date. All planting dates had a test weight that met the industry standard. Oil content was highest in the earliest-planted sunflower seeds (28.7%), but not significantly higher than sunflower oil content from the second or third planting dates. The overall oil yield was highest in sunflowers planted on 7-June or 25-May.

**Table 4. Effect of planting date on harvest yields and quality.**

Planting date	Moisture at harvest	Seed yield	Test weight	Insect damage to seed	Oil content	Oil yield	
	%	lbs / acre	lbs / bu	%	%	lbs / acre	gal / acre
5/25/2011	13.0	1410	26.3	<b>15.3</b>	<b>28.7</b>	386*	50.5*
6/1/2011	16.0	1150	25.8	17.2	19.6	225	29.4
6/7/2011	<b>21.0*</b>	<b>1540</b>	<b>27.8*</b>	19.8	27.1	<b>418*</b>	<b>54.8*</b>
LSD (0.10)	2.8	NS	1.2	NS	NS	122	15.9
Trial Mean	16.7	1360	26.7	17.4	25.2	343	44.9

\* Treatments indicated with an asterisk did not perform significantly lower than the top-performing treatment in a particular column. NS – No significant difference was determined between treatments.

## DISCUSSION

In this trial, populations of sunflowers planted on the first and last planting dates (May 25 and June 7) were lower than the recommended 24,000-28,000 plants per acre. The lowest establishment rates (64% and 53%) were in the earliest and latest planting date, respectively, and may have been due to weather events surrounding the plantings. In the week of the first planting date (May 25), northern Vermont had 5.67 inches of precipitation above normal, according to USDA weather data from Burlington, VT (30-year average from 1961-1990). The week of the second planting date (June 1), which had the highest emergence rate, also had above average rainfall, but slightly lower than the observed increase one week earlier (3.92 inches above normal). The final planting date (June 7) fell within a week that had 7.62 inches of rainfall, or 4.66 inches greater than normal. While the entire spring planting season had high precipitation (for example, the five weeks between May 16 and June 19 included 23.45 inches greater than historical averages), significant rainfall events occurred during the weeks of the first and last planting dates, possibly accounting for lower emergence rates and eventual populations.

Insect and bird damage likely is correlated to the life cycles of the major pests for sunflowers. Banded sunflower moth (*Cochylis hospes*), which is a major insect pest in sunflowers in northern states, feeds on sunflower seeds and reduces kernel weight and overall yield. The adult moths are generally prevalent from mid-July to mid-August, and lay eggs on sunflower bracts. These eggs hatch and larvae burrow into sunflower heads between mid-July and mid-September, feeding on kernels and wreaking havoc. Though not significantly different, the levels of insect damage (exit holes distinctive to sunflower moth damage) were lowest in the sunflowers planted earliest. This suggests that the sunflowers planted on May 25 this year, which had, for the most part, bloomed by July 27, may have already reached physiological maturity when pests such as banded sunflower moths became most active, and the younger plants had more desirable seed heads at the time. Damage to seedheads is most severe when the buds have just begun to elongate but have not yet opened.

Bird damage was significantly greatest (23.5%) among the sunflowers planted earliest, probably because they had already formed full, palatable seeds when migrating flocks first began to pass through Alburgh, VT. The most destructive birds in the northeast, American goldfinches (*Spinus tristis*) and red-winged blackbirds (*Agelaius phoeniceus*), migrate in the fall and can cause great amounts of damage while traveling in large flocks. Blackbirds have been known to pass through northern Vermont in the end of August as well, which may account for the higher percentage of bird damage in the earliest sunflowers, which had bloomed and begun to form seed by August, when the birds were storing up fat reserves for

long flights. The latest sunflowers suffered bird damage of 6.6%, perhaps because they developed palatable seeds between the two large waves of bird flocks.

It is not surprising that the sunflowers planted on the latest planting date had the highest moisture levels, since they had not had as much time in the field to mature and dry. While seed yields were not significantly different among planting dates, oil yields did differ according to the timing of spring planting. Sunflowers planted on the third planting date of the study (June 7) yielded the highest oil quantities (418 lbs or 54.8 gal per acre). Sunflowers planted on the first date (May 25) did not have significantly lower oil yields (386 lbs or 50.5 gal per acre).

The two varieties in the study, having differing relative maturities, have varying oil yields depending on the timing of planting. Because the shorter-season variety (Croplan 306, RM 87) had a lower oil yield than Syngenta 7120 (RM 95) on the earliest planting date, it is reasonable to conclude that the length of time between planting and harvesting was too great for that variety. Because it takes the variety 306 approximately 87 days to reach physiological maturity and the majority of all the plots were harvested on the same day (September 26, which was 124 days after the first planting date), the sunflowers were likely harvested past the ideal date. This result points to the importance of harvesting sunflowers according to their suggested and observed relative maturities. While seed yields may not vary across varieties by planting date, oil yields are not always directly related, due to differing oil contents under varying conditions.

While a general rule of thumb suggests that sunflowers should be planted as soon as possible after the spring soils have warmed to at least 50-55°F, 2011 was a very unusual year for weather. Wet soil conditions postponed spring planting for many Vermont growers, and Tropical Storm Irene, which hit Vermont in late August of 2011 with heavy rainfall and catastrophic winds, caused damage and difficulty in harvesting for many farmers. Especially with this anomalous weather, it is important to bear in mind that the results discussed here represent only one year's worth of data, and are specific to one location in northern Vermont. Decisions should not be based solely on this information, but rather a compilation of research across varying seasons and locations should be considered.

## ACKNOWLEDGEMENTS

UVM Extension would like to thank Roger Rainville and the staff at Borderview Farm for their generous help with this research trial, as well as the Vermont Sustainable Jobs Fund and USDA Northeast SARE (LNE09-282) for funding this research. We would also like to thank Katie Blair, Chantel Cline, Amber Domina, Laura Madden, Susan Monahan, and Amanda Gervais for their assistance with data collection. This information is presented with the understanding that no product discrimination is intended and no endorsement of any product mentioned, nor criticism of unnamed products, is implied.

*UVM Extension helps individuals and communities put research-based knowledge to work.*

Issued in furtherance of Cooperative Extension work. Acts of May 8 and June 30, 1914, in cooperation with United States Department of Agriculture, University of Vermont Extension, Burlington, Vermont. University of Vermont Extension, and U.S. Department of Agriculture, cooperating, offers education and employment to everyone without regard to race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or familial status

