

Energy and Economic Returns by Crop Rotation*

ith rising energy costs, input costs, and variable grain prices, adjusting crop rotations could be a profitable change for some farm operations. Using research from the ISU Extension and Outreach Marsden Research farm, the rotational effects on energy usage and economic returns were compared by two, three, and four year rotations. The 2-year rotation included GMO Corn and soybeans (C-Sb), the 3 and 4-year rotations included non-GMO Corn, Soybeans, and Oats, with the addition of a 4th year alfalfa crop in the 4-year rotation. (C-Sb-O and C-SB-O/A-A). The energy analysis focuses on uses of energy that have a price associated with them. To successfully compare across rotations, a line was drawn to keep energy use contained to producing the crop. It does not take into account solar energy, or energy consumption outside the "farm gate." Storage, hauling, and handling past initial removal of the crop are not considered. These have an impact on overall energy usage and economic returns but can vary widely and would not make equal comparisons across the three rotations.

This study looks at the period from 2006-2011. A previous study by Cruse, Liebman, Raman, and Wiedenhoeft takes a similar approach for 2003-2008. One major change between the two was a switch from triticale to oats for the small grain rotation. Figure 1 shows average yields for corn and soybeans for the three rotations. Yield gains were present for the 3 and 4 year rotations that can be associated with the agronomic benefits of the crop rotations and management practices used.

Energy Study

Energy use was split into five categories. Within each category,

the BTU/units were applied to the actual inputs and field operations from the research farm field notes. The five categories included: seed, grain drying, field operations, pesticides, and fertilizer.

There is limited information on the energy production of seeds for clover, oats, and alfalfa. Previous research (Shapouri, Cruse) use one factor for corn seed production and another for all other seeds. A similar method was used in this study as well. Factors used were 6,320 BTU/lb for corn, and 1,333 BTU/lb for other seeds.

Field operations were categorized into preharvest and harvest operations, including hauling grain.

This study looks at the energy usage for grain drying in a category of its own, in order to show the high energy requirements that drying can capture. The average reduction in moisture was 3.9% across the span of the study. This was multiplied by the factor of 1,620 BTU/point/bushel dried,Quick file giving a total of 6,320 BTU per bushel. This was then applied to the average yield for each rotation.



Figure 1. Average Yields by Crop and Rotation, 2006-2011

Ann Johanns, extension program specialist aholste@iastate.edu Craig Chase, local food and farm program coordinator Matt Leibman, H.A. Wallace Chair for Sustainable Agriculture Fuel usage for every field operation came from the publication, "Fuel Usage by Field Operation," (www.extension. iastate.edu/agdm/crops/pdf/a3-27.pdf). For operations not specifically listed, the closest equivalent was used as a substitute. The energy values and sources are listed in Table 1.

Fertilizer applied was averaged across all years to give a long term picture of energy use. Application was based on need, so varied year to year. Amounts used were taken from the provided field notes.

Though minor, nitrogen as an adjuvant was included in the fertilizer category.

The energy usage of herbicide and insecticide applications were based on active ingredients in the amounts applied. On the 3 and 4-year rotations, application was done by banding, thus reducing the overall rates.

Results

The rotational results were similar to previous studies (Cruse, et. al), though at a slightly lower rate. This can be attributed to several adjustments done in the later years of the study, including reduced fertilizer application and improved pesticide efficiencies, both in the product and method of application. Another change was the switch to oats from triticale.

Table 1. Energy Values Used							
Input	Energy Value Un	its	Source				
Seed – Corn	6,320 B	TU/lb	Grabowski				
All other seed	1,333 B	TU/lb	Sheehan				
Ν	24,500 B	TU/lb	Shapouri et al. (2004)				
Р	4,000 B	TU/lb	Shapouri et al. (2004)				
К	3,000 B	TU/lb	Shapouri et al. (2004)				
Herbicides	101,034 B	TU/lb	Bhat et al.				
, Insecticides	113,932 B	TU/lb	Bhat et al.				
Grain Drying	6,320 B	TU/bu	Grabowski				
Diesel	138,690 B	TU/gallon	EIA				

Figure 2 shows average energy inputs over the 6-year period, with the 2-year rotation clearly demanding the most energy. From 2006-2011, the three year rotation showed the least amount of energy usage based on the parameters of the study. Figure 3 illustrates in more detail where energy is expended. The highest category for energy usage was fertilizer for the two-year rotation, whereas the 3 and 4-year rotations show the most energy usage in field operations. Table 2 illustrates the details of percent of energy usage by category as indicated in Figures 2 and 3. In the 2-year rotation, just over 60% of energy expenditures came from fertilizer. In the 3 and 4-year rotations, the leading category of field operations carried 44 and 47%hhh, respectively, of the energy demands.



Figure 2. Average Energy Inputs by



Table 2. Percent of Energy Use byCategory

	GMO	Non-G	GMO
	C-SB	C-SB-O	C-SB-O-A
Seed	3%	8%	6%
Grain Drying	17%	30%	21%
Field Operations	14%	44%	47%
Pesticides	5%	1%	1%
Fertilizer	61%	17%	25%

Energy use was looked at by crop as well as by rotation. In Figure 4, the two-year corn crop leads as the most energy demanding, much higher than any other crop. The two-year corn crop utilized more than twice the energy of any other crop at 5.83 million BTU, primarily due to increased herbicide and fertilizer usage. Table 3 shows the 3 and 4-year corn rotations come fairly close to one another at 2.45 and 2.63 million BTU each. The reason for the difference in those rotations can mostly be attributed to the drying costs associated with the slight higher corn yields in the 4-year rotation.

Table 3. Total BTU by Crop

	2yr	3yr	4yr
Corn	5.83	2.45	2.63
Soybeans	1.23	1.00	1.13
Oats		0.77	1.09
Alfalfa (2nd yr)			1.14
Total Average BTU by Rotation	3.53	1.41	1.50

Figure 4. Energy Input per Crop for Individual Rotations





Figure 5. Energy Usage by Rotation in Diesel Fuel Equivalents, 2006-2011

Another way to look at energy usage is in diesel fuel equivalents. This is shown in Figure 5. Diesel fuel equivalents were found by taking the total BTU/rotation divided by the BTU/gallon of diesel fuel. This represents the energy consumption in an easily recognizable form, even though not all energy usage was associated with diesel fuel. The two year rotation uses the equivalent of 25.43 gallons of diesel fuel per acre. The 3 and 4-year are both just over 10 gallons per acre.

Economic Returns

Along with energy usage, this study also compares economic returns for the three rotations. Using data from the annual publication for estimated costs of crop production for that year, the costs for production were applied to the various cost components (Duffy, et. al). Herbicide and Insecticide prices were taken from annual reports from North Dakota State University and the University of Nebraska. Annual grain prices came from the USDA National Ag Statistics Services, Iowa office. No government payments or other income were included in the study. With differences in field operations, fertilizer, and reduced pesticides, the three and four year rotations have the ability to compete with the two-year rotation in profitability as well as energy requirements.

Energy Usage by Rotation in Diesel Fuel Equivalents						
	2yr	Зуr	4yr			
Diesel Fuel Equivalents (gal/acre)	25.43	10.16	10.80			

Figure 6 shows the average return to management by crop and rotation. Figure 7 shows the average returns for the three rotations to land, labor and management, land and management, and management. The first of these categories shows the returns if the costs for land, labor, and management are not included. The second takes into account the cost for labor, and the last includes all costs and what remains are the returns to management. Rotational effects of increased yields and lower input costs for the non-GMO crops make the three year rotation result in the highest returns for this study. Table 4 gives a detailed summary of the cost and returns by crop and rotation and shows the three year rotation having an average Return to Management of \$194.03; and the two-year and fouryear rotations having positive returns of \$187.92 and \$170.97, respectively.







Table 4. Revenue and Returns by Crop and Rotation

	Yields	Prices	Gross Revenue	Production Cost	Return LL&M	Return L&M	Return Mgmt
C-Sb							U
Corn	193.73	\$4.35	\$841.60	\$377.27	\$464.33	\$457.25	\$259.09
Soybeans	50.27	9.95	503.82	180.76	323.06 \$393.70	314.92 \$386.08	116.75 \$187.92
C-Sb-O					<i></i>	<i></i>	÷
Corn	198.80	\$4.35	\$865.19	\$255.22	\$609.97	\$592.73	\$394.57
Sovbeans	54.73	9.95	549.31	158.01	392.00	380.72	182.56
Oats	97.92	2.69	256.35	129.54	212.04	203.13	4.97
Oat Straw	1.07	79.17	85.23				
					\$404.67	\$392.20	\$194.03
C-Sb-O/A-A						·	
Corn	202.43	\$4.35	\$878.09	\$275.72	\$602.37	\$584.75	\$386.59
Soybeans	56.93	9.95	571.12	175.97	395.15	383.91	185.74
Oats	101.58	2.69	267.15	215.37	218.05	203.72	5.55
Oat Straw	1.00	79.17	79.74				
Alfalfa (1st yr)	0.74	119.47	86.52				
Alfalfa (2nd yr)	3.97	119.47	470.21	145.08	325.13 \$385.17	304.16 \$369.14	106.00 \$170.97

Nutrient Pricing

A portion of the cost savings in the three and four year rotations was due to applying manure from an available livestock operation to all corn acres in the rotation. In the initial study, the only cost associated with the manure is the cost to apply. Another way to phrase this is that manure was viewed as a "waste product" that needs to be disposed of in a feasible manner.

Another second method was also applied to see how it would affect the profitability of the rotations. This involved valuing the manure based on its nutrient value. The concept behind this method was to show the cost that would have occurred had an equivalent amount of fertilizer been purchased commercially.

Manure was analyzed each year for nutrient content and these rates were used along with the cost that would have been assessed had those nutrients been purchased. Nutrient prices used were from the ISU publication for Estimated Costs of Crop Production. A third option in comparing the rotations might be to put a flat rate per ton or load on the manure rather than breaking the cost down for each nutrient. The energy usage of the animals in producing the manure is not considered. This could also be taken into consideration as far as number of animals, manure nutrient content, etc. Figures 8 and 9 show the returns to land, labor, and management by crop and rotation. Table 5 gives the economic data with manure priced at its nutrient value. Applying this process shows the benefits of having manure readily available and that it is a major factor in the profitability of the non-GMO rotations in this study.

Summary

This publication has focused on the energy use and economic returns of three different crop rotations. The choice of which rotation to choose is dependent on many factors. When considering profitability and energy consumption, including a third or fourth crop may be a viable option for some operations. Other benefits might include an outlet for excess manure, reduced erosion, increased soil health and pest management.

Table 5. Revenue and Returns by Crop and Rotation, with manure priced by nutrient value							
	Yields	Prices	Gross Revenue	Production Cost	Return LL&M	Return L&M	Return Mgmt
C-Sb							U
Corn	193.73	4.35	\$841.60	377.27	464.33	\$457.34	\$259.18
Soybeans	50.27	9.95	\$503.82	180.76	323.06 393.70	\$314.92 386.13	\$116.75 187.96
C-Sb-O							
Corn	198.80	4.35	\$865.19	336.72	528.47	\$511.24	\$313.07
Sovbeans	54.73	9.95	\$549.31	157.32	392.00	\$380.72	\$182.56
Oats	97.92	2.69	\$256.35	128.56	213.01	\$204.42	\$6.26
Oat Straw	1.07	79.17	\$85.23			r -	· · ·
			·		377.83	365.46	167.29
C-Sb-O/A-A							
Corn	202.43	4.35	\$878.09	357.22	520.87	\$503.26	\$305.09
Soybeans	56.93	9.95	\$571.12	175.97	395.15	\$383.91	\$185.74
Oats	101.58	2.69	\$267.15	215.12	218.30	\$203.97	\$5.80
Oat Straw	1.00	79.17	\$79.74			·	
Alfalfa (1st vr)	0.74	119.47	\$86.52				
Alfalfa (2nd yr)	3.97	119.47	\$470.21	144.11	326.10 365.11	305.22 349.09	107.06 150.92

Figure 8. Average Return to Management by Crop and Rotation





Figure 9. Average Returns to Land, Labor, and Management by Crop Rotation



References

Chase, C., Delate, K., Liebman, M., Leibold, K. 2008. Economic Analysis of Three Iowa Rotations. Iowa State University Extension and Outreach, Bulletin PMR 1001. http://www.extension.iastate. edu/Publications/PMR1001.pdf

Cruse, M.J., Liebman, M., Raman, D.R., Wiedenhoeft, M. 2010. Fossil Energy Use in Conventional and Low-External-Input Cropping Systems. Agronomy Journal, 102(3): 934-941. Duffy, M., Smith, D. 2006-2011. "Estimated costs of crop production in Iowa." Ames, Iowa: Iowa State University, Extension and Outreach, FM-1712.

Graboski, M.S. 2002. Fossil Energy Use in the Manufacture of Corn Ethanol. National Corn Growers Association, St. Louis, MO. On-line at: http://www.oregon.gov/ENERGY/RENEW/ Biomass/docs/FORUM/FossilEnergyUse.pdf?ga=t

Page 8

Sheehan, J., Camobreco, V., Duffield, J., Graboski, M. & Shapouri, H. 1998. Life Cycle Inventory of Biodiesel and Petroleum Diesel for Use in an Urban Bus (Natl. Renewable Energy Lab., Golden, CO), NREL Publ. No. SR-580-24089. On-line at: http:// www.nrel.gov/docs/legosti/fy98/24089.pdf

USDA. 2006-2011. Annual Grain Prices. Des Moines, IA: United States Department of Agriculture, National Ag Statistics Services, Iowa Office.

Guide for Weed Management. 2006-2011. University of Nebraska Extension. Lincoln, NE.

ND Weed Control Guide. 2006-2011. North Dakota State University. Fargo, ND. On-line at: http://www. ndsu.edu/weeds/weed-control-guides/nd-weedcontrol-guide-1/

* Funds for this project were provided by the Leopold Center for Sustainable Agriculture and Practical Farmers of Iowa.

. . . and justice for all

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Many materials can be made available in alternative formats for ADA clients. To file a complaint of discrimination, write USDA, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and July 30, 1914, in cooperation with the U.S. Department of Agriculture. Cathann A. Kress, director, Cooperative Extension Service, Iowa State University of Science and Technology, Ames, Iowa.