

Table 1. Proportion of native legume seed recruited to seedling stage for five species of native legumes measured at three sampling periods.

a. Desmodium canadense (showy tick trefoil)

	Fall 2009			June 2010			Fall 2010		
Diversity	Cont.	Graze	Mow	Cont.	Graze	Mow	Cont.	Graze	Mow
SW	0.04	0.19	0.08	0.00	0.08	0.12	0.05	0.02	0.02
LD	0.07	0.08	0.14	0.07	0.24	0.13	0.02	0.01	0.02
HD	0.04	0.12	0.13	0.28	0.26	0.07	0.02	0.07	0.05

b. Desmanthus illinoensis (Illinois bundle-flower)

	Fall 2009			June 2010			Fall 2010		
Diversity	Cont.	Graze	Mow	Cont.	Graze	Mow	Cont.	Graze	Mow
SW	0.02	0.04	0.03	0.07	0.07	0.07	0.00	0.01	0.01
LD	0.04	0.04	0.04	0.03	0.13	0.07	0.01	0.00	0.00
HD	0.02	0.08	0.11	0.03	0.10	0.15	0.00	0.01	0.01

c. Dalea purpurea (purple prairie-clover)

	Fall 2009			June 2010			Fall 2010		
Diversity	Cont.	Graze	Mow	Cont.	Graze	Mow	Cont.	Graze	Mow
SW	0.01	0.06	0.02	0.00	0.03	0.03	0.00	0.00	0.00
LD	0.00	0.03	0.05	0.00	0.00	0.00	0.00	0.00	0.00
HD	0.01	0.06	0.04	0.00	0.00	0.00	0.00	0.00	0.00

d. Lespedeza capitata (round-headed bush-clover)

	Fall 2009			June 2010			Fall 2010		
Diversity	Cont.	Graze	Mow	Cont.	Graze	Mow	Cont.	Graze	Mow
SW	0.01	0.12	0.05	0.00	0.03	0.00	0.00	0.00	0.00
LD	0.04	0.09	0.05	0.00	0.00	0.00	0.00	0.00	0.00
HD	0.01	0.14	0.08	0.00	0.00	0.00	0.00	0.00	0.00

e. Amorpha canescens (lead-plant)

	Fall 2009			June 2010			Fall 2010		
Diversity	Cont.	Graze	Mow	Cont.	Graze	Mow	Cont.	Graze	Mow
SW	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00
LD	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
HD	0.00	0.01	0.00	0.00	0.03	0.00	0.00	0.00	0.00