

Table 1 Sampling summary by grass species from three spring harvests.

| Species in Mix | 2011 ^a | | 2012 ^b | | 2013 ^c | |
|-------------------|-------------------|------------------------|-------------------|------------------------|-------------------|------------------------|
| | n | % Grass (Mean ± SD) | n | % Grass (Mean ± SD) | n | % Grass (Mean ± SD) |
| Timothy | 163 | 42.3 ± 16.7 | 59 | 47.8 ± 12.8 | 52 | 57.7 ± 8.8 |
| Reed canarygrass | 166 | 39.6 ± 16.0 | 60 | 53.6 ± 18.0 | 35 | 54.2 ± 9.1 |
| Orchardgrass | 191 | 58.1 ± 14.4 | 60 | 74.5 ± 16.9 | 12 | 64.7 ± 8.3 |
| Quackgrass | 55 | 62.2 ± 12.6 | | | | |
| Tall Fescue | | | | | 50 | 41.4 ± 23.3 |
| Smooth bromegrass | | | | | 38 | 60.5 ± 8.5 |
| Pure Alfalfa | 5 | 0 | | | | |

^a 2011 images were acquired with a single digital camera: Canon PowerShot A3100 IS.

^b 2012 images were acquired with four digital cameras including: Canon PowerShot A3100 IS, iPhone 4, Olympus FE230/X790, and Nikon Coolpix S6200.

^c 2013 images were acquired with three digital cameras including: iPhone 4, Nikon Coolpix S6200, and Nikon Coolpix E7600. Two samples were missing from the Nikon Coolpix S6200.