Table 5. Average (n=4) total nutrient element content of soil (± standard deviation) prior to seaweed addition (October 2011) and at the end of the growing season (September 2012). Within columns, values with the same letter are not significantly different.

Fertilizer	K		Ca		Fe		Mn	
treatment								
	g/kg dry soil							
	Oct	Sep	Oct	Sep	Oct	Sep	Oct	Sep
LDS	14.8^{a}	15.0^{a}	8.08^{a}	7.93^{a}	19.7^{a}	18.6^{a}	0.330^{a}	0.351^{a}
	± 0.49	± 0.23	± 0.31	± 0.21	± 1.3	± 0.15	± 0.0025	± 0.0019
HDS	15.0^{a}	15.2^{a}	8.02^{a}	7.96^{a}	19.3 ^a	18.5 ^a	0.314^{a}	0.302^{a}
	± 0.59	± 0.31	± 0.20	± 0.39	± 0.47	± 0.23	± 0.0026	± 0.0024
PFF	15.3 ^a	15.0^{a}	8.21^{a}	7.94^{a}	18.8 ^a	19.0^{a}	0.314^{a}	0.351^{a}
	± 0.15	± 0.20	± 0.39	± 0.007	± 1.3	± 0.30	± 0.0057	± 0.0044