

Table 4. Average forage height (Least-squares means) before and after grazing different treatments during the 2012-2013 cool-season growing period (De. or Jan.-Apr.), Phenix City, AL, USA.

Treatment	Forage height			
	Grass		Legume	
	Before	After	Before	After
	-----cm-----			
ArrowleafRye [†]	21 ^{‡b} _x	7 ^b _y	11 ^c _x	4 ^b _y
BerseemRye	21 ^b _x	8 ^b _y	11 ^c _x	5 ^b _y
CrimsonRye	22 ^b _x	7 ^b _y	15 ^b _x	4 ^b _y
PeasRye	25 ^a _x	10 ^a _y	25 ^a _x	18 ^a _y
Marshall ryegrass	22 ^b _x	8 ^b _y	-	-
VetchRye	21 ^b	8 ^b	10 ^d _x	3 ^b _y

[†] ArrowleafRye = Arrowleaf clover (*Trifolium vesiculosum*)-Marshall ryegrass (*Lolium multiflorum*)

BerseemRye = Berseem clover (*Trifolium alexandrinum*)-Marshall ryegrass

CrimsonRye = Crimson clover (*Trifolium incarnatum*)-Marshall ryegrass

PeasRye = Winter peas (*Pisum sativum*)-Marshall ryegrass

VetchRye = Hairy vetch (*Vicia villosa*)-Marshall ryegrass

[‡] Least-squares means for forage height within a column with different superscripts, and within a row under grass or legume category with different subscripts are different (P < 0.05).