

Literature Cited.

- Barnett, K. 2012. Wisconsin Fresh Market Vegetable Budgets: Pumpkins. <http://cdp.wisc.edu/Freshmarket.htm>
- Bianchi, F.J., Booij, C.J. and Tscharntke T. 2006. Sustainable pest regulation in agricultural landscapes: a review on landscape composition, biodiversity and natural pest control. *Proceedings of the Royal Society of London Series B* 273:1715-1727.
- Biesmeijer, J., Roberts, S., Reemer, M, Ohlemüller, R., Edwards, M., Schaffers, A., Potts, S., Kleukers, S., Thomas, C., Settele, J., and W. Kunin. 2006. Parallel Declines in Pollinators and Insect-Pollinated Plants in Britain and the Netherlands. *Science* 313:351-354.
- Blaauw, B. R., and Isaacs, R. 2012. Larger wildflower plantings increase natural enemy density, diversity, and biological control of sentinel prey, without increasing herbivore density. *Ecological Entomology* 37(5), 386–394. doi:10.1111/j.1365-2311.2012.01376.x
- Carvell, C., D. Roy, S. Smart, R. Pywell, C. Preston, and D. Goulson. 2006. Declines in forage availability for bumblebees at a national scale. *Biological Conservation* 132:481-489.
- Fiedler, A. K. and D. A. Landis. 2007a. Attractiveness of Michigan native plants to arthropod natural enemies and herbivores. *Environmental Entomology* 36:751-765.
- Fiedler, A. K. and D. A. Landis. 2007b. Plant characteristics associated with natural enemy abundance at Michigan native plants. *Environmental Entomology* 36:878-886.
- Fiedler, A. K. and D. A. Landis. 2008. Maximizing ecosystem services from conservation biological control: the role of habitat management. *Biological Control* 45:254-271.
- Isaacs, R., and Kirk, A. K. 2010. Pollination services provided to small and large highbush blueberry fields by wild and managed bees. *Journal of Applied Ecology*, 47(4), 841–849. doi:10.1111/j.1365-2664.2010.01823.x
- Gajc-Wolska, J., Kowalczyk, K., and Jarosław, M. 2011. Efficiency of cucumber (*Cucumis sativus* L.) pollination by bumblebees (*Bombus terrestris*). *Acta Sci. Pol., Hortorum Cultus*, 10(1), 159–169.
- Goverde, M., Schweizer, K., Baur, B., and Erhardt, A. 2002. Small-scale habitat fragmentation effects on pollinator behaviour: experimental evidence from the bumblebee *Bombus veteranus* on calcareous grasslands. *Biological Conservation*, 104(3), 293–299. doi:10.1016/S0006-3207(01)00194-X
- Kleijn, D. and W. Sutherland. 2003. How effective are European agri-environment schemes in

conserving and promoting biodiversity? *Journal of Applied Ecology* 40:947-969.
enemies of arthropod pests in agriculture. *Annual Review of Entomology* 45:175-201.

Long, R. F., A. Corbett, C. Lamb, C., Reberg-Horton, Chandler, J., and Stimmann, M. 1998.
Beneficial insects move from flowering plants to nearby crops. *California Agriculture* 52:23-26.

Mitchell, T.B. (1962) Bees of the eastern United States. II. Technical bulletin (North Carolina
Agricultural Experiment Station), 152, 1-557. [Megachilidae, Anthophoridae, Apidae s.s.]

Petersen, J. D., Reiners, S., and Nault, B.A. 2013. Pollination services provided by bees in
pumpkin fields supplemented with either *Apis mellifera* or *Bombus impatiens* or not
supplemented. *PloS One*, 8(7), e69819. doi:10.1371/journal.pone.0069819

Rebek, E. J., Sadof, C. S., and Hanks, L. M. 2005. Manipulating the abundance of natural
enemies in ornamental landscapes with floral resource plants. *Biological Control* 33:203-
216.

Smith, A. A., Bentley, M., Reynolds, H. L., and Smith, A. A. 2013. Wild Bees Visiting
Cucumber on Midwestern U . S . Organic Farms Benefit from Near-Farm Semi-Natural Areas
Wild Bees Visiting Cucumber on Midwestern U . S . Organic Farms Benefit From Near-Farm
Semi-Natural Areas. *Journal of Economic Entomology*, 106(1), 97–106.

Sotherton, N. W. 1998. Land use changes and the decline of farmland wildlife: An appraisal of
the set-aside approach. *Biological Conservation* 83:259-268.

Stanghellini, M. S., Ambrose, J. T., and Schultheis, J. R. 1997. The Effects of Honey Bee and
Bumblebee Pollination on Fruit Set and Abortion of Cucumber and Watermelon. *American Bee
Journal*, 137(5), 386–391.

Tuell, J. K., Fiedler A. K., D. A. Landis, and Isaacs R. 2008. Visitation by wild and managed
bees (Hymenoptera: Apoidea) to Eastern U.S. native plants for use in conservation programs.
Environmental Entomology 37:707-718.

Van Lenteren, J. C. 2011. The state of commercial augmentative biological control: plenty of
natural enemies, but a frustrating lack of uptake. *BioControl*, 57(1), 1–20. doi:10.1007/s10526-
011-9395-1

Walton, N. J., and Isaacs, R. 2011. Influence of native flowering plant strips on natural enemies
and herbivores in adjacent blueberry fields. *Environmental Entomology*, 40(3), 697–705.
doi:10.1603/EN10288

Wanner, H., Gu, H., and Dorn, S. 2006. Nutritional value of floral nectar sources for flight in the
parasitoid wasp, *Cotesia glomerata*. *Physiological Entomology* 31:127-133.

Williams, N. M., Crone, E. E., Roulston, T. H., Minckley, R. L., Packer, L., and Potts, S. G. 2010. Ecological and life-history traits predict bee species responses to environmental disturbances. *Biological Conservation*, 143(10), 2280–2291. doi:10.1016/j.biocon.2010.03.024

Winfree, R., Gross, B. J., and Kremen, C. 2011. Valuing pollination services to agriculture. *Ecological Economics*, 71, 80–88. doi:10.1016/j.ecolecon.2011.08.001

Vegetables 2013 Summary. 2014. United States Department of Agriculture, National Agricultural Statistics Service 1-82.

United States Standards for Grades of Cucumbers. 1997. United States Department of Agriculture, Agricultural Marketing