

Building Soil Organic Matter

Sam Corcoran, Alex Wu, Ruichen Wang, Masoud Hashemi

Soil Organic Matter (SOM) is the fraction of the soil that consists of plant and animal residues in various stages of decomposition. It contains organic carbon and nitrogen. For soil microorganisms, such as fungi and bacteria, the carbon is used as a source of energy whereas nitrogen is used as a source of protein. In healthy soils, the vast majority of soil microorganisms are beneficial or neutral and can help prevent organisms such as plant pathogens from becoming dominant.

SOM is important because it improves many physical, chemical, and biological characteristics of soil. When SOM is well decomposed, it enhances biological activities in the soil, improves soil structure by increasing aggregation and aggregation stability, and slowly releases nitrogen and phosphorus, reducing the need for synthetic fertility amendments. Additional SOM benefits include increased ability of the soil to hold water, reduced erosion, stabilized (buffered) pH, and increased availability of micronutrients. A loss of SOM can lead to negative effects such as soil erosion, loss of fertility, compaction, and land degradation. Because SOM contains an average of 58% carbon by volume, building soil organic matter significantly contributes to carbon sequestration. Therefore, farming practices that increase amounts of SOM, or reduce its loss, are ideal.

Building SOM is a very slow process that begins with photosynthesis. During photosynthesis, plants take carbon from the atmosphere and form sugars. As much as 40-44% of these sugars are then released from plant roots as root exudates. These sugars help feed the soil microbial communities, and the sugars become a part of the microbes' bodies. In return, these microbes can increase soil fertility and health by suppressing plant diseases, delivering nutrients, and increasing water use efficiency. Likewise, when plants die or when manure is added to the soil, the microbes can decompose these materials. Thus, soil organic matter is found in three states:

1. Living: microbes and other soil biology
2. Dead: plant residues and manures added to the soil
3. Very Dead: this is what we call humus. The living SOM digests the dead SOM to make the very dead SOM.

When we want to increase SOM, we must set reasonable expectations about how quickly this will occur. Let us suppose that you want to increase your SOM by 1%. First, we must know that one acre of soil to a depth of 6 inches (acre furrow slice) weighs approximately 2,000,000 lbs. One percent of 2,000,000 is 20,000 lbs. of SOM.

It is necessary to convert SOM to carbon to continue our equation. To convert between soil organic matter to soil organic carbon, the Van Bemmelen conversion factor of 0.58 is used. This number assumes that all SOM is approximately 58% carbon. There is variation in this number, but this conversion factor is widely used and is an accepted estimate. With this conversion factor, 20,000 lbs. of SOM equals 11,600 lbs. of carbon. This means that approximately 11,600 lbs. of carbon from plant or manure residues is required to increase soil organic matter by 1%.

However, 11,600 lbs. of carbon means a lot of plant matter or manure is needed. For example, the carbon content by mass of tall fescue, a common pasture and hay grass, is 47%. Thus, to increase SOM by 1%, you would need to add 24,680 lbs. of fescue dry matter ($0.47x=11,600$, solve for x).

Yet, in reality, it is estimated that 10-20% of plant residue converts into SOM. The remaining components are lost back to the environment in the breakdown process. Adding this value into the calculations, you would need 123,400 to 246,800 lbs. of fescue dry matter to increase SOM by 1% ($0.1x=24680$ to $0.2x=24680$, solve for x). In a productive season of well managed pasture or hay, tall fescue may yield around 9000 lbs. of above ground dry biomass per acre. It would take years of biomass incorporation to reach the desired 1% SOM increase.

It is critical to point out that this calculation only considers the aboveground biomass of tall fescue. In reality, the roots of the fescue and the root exudates can also convert into SOM. In some crops, the root contribution can be nearly half of the total annual crop residues for that crop. In others, root contributions can even be 2.3 times the aboveground biomass amount. The root residues provide important inputs up to 8 inches deep in the soil to feed soil organisms. If we estimate that the roots contribute the same amount of carbon and SOM as the above ground biomass, it would take 6 years of fescue growth – with all the aboveground biomass being returned to the soil – to reach a 1% increase to SOM.

A series of similar calculations and estimates, based on field research, have led to the generalization that we can expect a 0.1% increase to SOM per year in a well-managed system. Yet, in-field variation when we take soil samples, coupled with the margin of error on laboratory analysis of SOM, means it is very difficult to reliably identify increased to SOM in the short term. Therefore, it is recommended that changes to SOM be measured after a period of 3-5 years for a reliable indicator. Large, sudden increases to SOM are best explained by errors, not real change in the soil.

Currently, more research is being conducted on how much root biomass and exudates convert to SOM. This will improve our estimates in the future. Plant roots have a different chemical composition than aboveground biomass, but have close association with soil microorganisms. Therefore, roots decompose at different rates compared to aboveground biomass like leaves.

Likewise, roots and their exudates are already in the soil, whereas aboveground biomass must work its way into the soil. This can of course be affected by the tillage method chosen to manage agricultural land.

Table 1: How many pounds of dry cover crop biomass is needed for 1% increase in SOM, assuming a 10-20% conversion rate. Roots are not included in these estimates.

Crop/Biomass Input	Carbon Content	lbs Dry Matter for 1% Increase in SOM
Annual rye/winter rye	42%	138,095-276,190
Tall fescue	47%	123,400-246,800
Vetch	38%	152,631-305,263
Vetch and rye	39%	148,718-297,436

Roots have long been undermeasured due simply to the difficulty of removing roots from the soil and washing them in order to estimate biomass. However, this work is expected to increase in the immediate future. In the meantime? Emphasize practices that build SOM, set reasonable expectations, and give it time.

This material supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture, through the Northeast Sustainable Agriculture Research and Education program under subaward number SNE20-004-MA.

References

- Burns, J. C., Chamblee, D. S., & Giesbrecht, F. G. (2002). Defoliation Intensity Effects on Season-Long Dry Matter Distribution and Nutritive Value of Tall Fescue. *Crop Science*, 42(4), 1274–1284. <https://doi.org/10.2135/cropsci2002.1274>
- Canarini, A., Kaiser, C., Merchant, A., Richter, A., & Wanek, W. (2019). Root Exudation of Primary Metabolites: Mechanisms and Their Roles in Plant Responses to Environmental Stimuli. *Frontiers in Plant Science*, 10. <https://www.frontiersin.org/articles/10.3389/fpls.2019.00157>
- Chenu, C., Angers, D. A., Barré, P., Derrien, D., Arrouays, D., & Balesdent, J. (2019). Increasing organic stocks in agricultural soils: Knowledge gaps and potential innovations. *Soil and Tillage Research*, 188, 41-52.
- Finney, D. M., White, C. M., & Kaye, J. P. (2016). Biomass Production and Carbon/Nitrogen Ratio Influence Ecosystem Services from Cover Crop Mixtures. *Agronomy Journal*, 108(1), 39–52.
- Fuhrmann, J. J., & Zuberer, D. A. (2021). Carbon Transformations and Soil Organic Matter Formation. *Principles and Applications of Soil Microbiology*, 327–361.
- Kamilova, F., Kravchenko, L. v, Shaposhnikov, A. I., Azarova, T., Makarova, N., & Lugtenberg, B. (2006). Organic Acids, Sugars, and L-Tryptophane in Exudates of Vegetables Growing on Stonewool and Their Effects on Activities of Rhizosphere Bacteria. *Molecular Plant-Microbe Interactions MPMI*, 19(3), 250–256. <https://doi.org/10.1094/MPMI>
- Kätterer, T., Bolinder, M. A., Andrén, O., Kirchmann, H., & Menichetti, L. (2011). Roots contribute more to refractory soil organic matter than above-ground crop residues, as revealed by a long-term field experiment. *Agriculture, Ecosystems and Environment*, 141(1–2), 184–192. <https://doi.org/10.1016/j.agee.2011.02.029>
- Kuzyakov, Y., & Cheng, W. (2001). Photosynthesis controls of rhizosphere respiration and organic matter decomposition. *Soil Biology and Biochemistry*, 33(14), 1915–1925. [https://doi.org/https://doi.org/10.1016/S0038-0717\(01\)00117-1](https://doi.org/https://doi.org/10.1016/S0038-0717(01)00117-1)
- Qian, Y. L., Fry, J. D., & Upham, W. S. (1997). Rooting and Drought Avoidance of Warm-Season Turfgrasses and Tall Fescue in Kansas. *Crop Science*, 37(3), 905–910. <https://doi.org/10.2135/cropsci1997.0011183X003700030034x>
- Roth, T., & Waite, J. (2021). Early Spring Carbon to Nitrogen Ratios of Cereal Rye Varieties. <https://www.ncei.noaa.gov/access/us-climate-normals/>

Sainju, U. M., Whitehead, W. F., & Singh, B. P. (2005). Biculture Legume–Cereal Cover Crops for Enhanced Biomass Yield and Carbon and Nitrogen. *Agronomy Journal*, 97(5), 1403–1412.
<https://doi.org/10.2134/agronj2004.0274>

USDA - Natural Resources Conservation Service Booneville Plant Materials Center. (2007). Dry-Matter Production of Eight Grass Species With Three Levels of Poultry Litter. *Plant Materials Technical Note*.
https://www.nrcs.usda.gov/Internet/FSE_PLANTMATERIALS/publications/arpmctn7593.pdf