

Table 1. Descriptions of materials, rates, and costs of materials used in the field study.

	Product	Active Ingredient	Interval	Rate	Cost (\$) / Tunnel/Season	Cost (\$) / Tunnel/Application
1	Untreated	n/a				
2	Double Nickel LC	<i>Bacillus amyloliquefaciens</i> Strain D747	2 wks	1.5 qt/100 gall	114.13	9.51
3	Rootshield Plus	<i>Trichoderma harzianum</i> Rifai strain T-22 (1.15%); <i>Trichoderma virens</i> strain G-41 (0.61%)	6 wks	8 oz/100 gall	74.82	18.71
4	Actinovate AG	<i>Streptomyces lydicus</i> WYEC 108 (0.0371%)	2 wks	9 oz/100 gall	245.39	20.45
5	Mycostop G	<i>Streptomyces griseoviridis</i> Strain K61 (35%)	4 wks	2 g/100 sq ft	2073.60	345.60
6	Oxidate	Hydrogen dioxide	1 wks	1:300	753.21	31.38
7	RootShield Plus, Oxidate	<i>Trichoderma harzianum</i> Rifai strain T-22 (1.15%), <i>Trichoderma virens</i> strain G-41 (0.61%); Hydrogen Dioxide	6 wks; 1 wks	8 oz/100 Gall; 1:300	828.03	50.09