

**Table 2.** Pest suppression following termination of a mustard cover crop is caused by the enzymatic hydrolysis of glucosinolates. In *Brassica juncea* — the mustard species in this project — the primary glucosinolate is sinigrin. This table presents sinigrin concentrations in *Brassica juncea* cv ‘rojo’ grown at four sites in southern New Mexico. Plant tissues were collected just before plant termination. The table also presents literature reports of sinigrin concentrations for *B. juncea* grown under field conditions.

Site	<i>Brassica juncea</i> cv ‘rojo’		Concentration ranges reported in literature				
	Green phenotype <sup>1</sup>	Purple phenotype	Bangarwa et al., 2011 <sup>2</sup>	Villata et al., 2016 <sup>3</sup>	Doheny-Adams et al., 2018 <sup>4</sup>	Kirkegaard and Sarwar, 1999 <sup>5</sup>	Ngala et al., 2015 <sup>6</sup>
	μmol g <sup>-1</sup> shoot tissue						
Columbus	43.9	42.5					
Deming	NA	NA					
Las Uvas	37.2	37.6	0.5 – 72.3	29 – 72	6 – 37	5 – 35	10 – 112.5
Leyendecker	45.4	42.9					

<sup>1</sup> *Brassica juncea* cv ‘rojo’ included green and purple phenotypes

<sup>2</sup> *Weed Science* 59:247-254

<sup>3</sup> *Journal of Agricultural Chemistry and Environment* 5:38-45

<sup>4</sup> *Journal of Agricultural and Food Chemistry* 66:5108-5116

<sup>5</sup> *Australian Journal of Agricultural Research* 50:315-324

<sup>6</sup> *Pest Management Science* 71:759-769