

Table 2. Annual bahiagrass production from multiple Florida locations under different fertility management.

Treatment	Year	Osceola	Orange	Columbia	Gulf
		----- (kg ha ⁻¹) -----			
Complete fertilizer	1	7,103	3,135	8,549	10,127
	2	7,448	8,558	7,162	7,325
	3	7,909	10,228	9,653	11,266
	6	7,998	NA	NA	NA
Class AA biosolids ²	1	6,599	3,219	4,744	9,878
	2	8,553	8,916	7,295	6,355
	3	6,885	9,944	8,044	10,305
	6	7,177	NA	NA	NA
Minus P fertilizer	1	7,320	3,647	7,317	8,785
	2	9,716	9,018	7,101	6,117
	3	9,230	9,701	9,404	10,504
	6	7,824	NA	NA	NA
Minus K fertilizer	1	6,080	3,778	4,681	7,161
	2	6,308	7,973	5,904	4,919
	3	5,192	8,476	6,872	5,584
	6	2,364	NA	NA	NA
Minus N fertilizer	1	3,008	3,199	3,230	6,422
	2	3,643	6,468	4,990	5,096
	3	2,368	4,831	4,408	5,905
	6	2,914	NA	NA	NA
No fertilizer (check)	1	2,848	3,241	2,052	4,571
	2	5,566	4,961	3,623	3,360
	3	2,113	3,027	2,291	5,550
	6	1,797	NA	NA	NA

²Class AA biosolids + KCl (0-0-60)