

**Table 3. The seasonal availability of the 10 most abundant wildflower species used in the agricultural areas of Pinefield Ecofarm (Hepzibah, Georgia). These species exhibited the best survivorship in the sandy soils of the coast plain in Eastern Georgia. Blueberry (*Vaccinium species*) and cucumber (*Cucumis sativus*) were the target agricultural crops. They are listed with their flowering times at the top of the table.**

xx = present

Plants	Common Name	Feb.	Mar.	Apr.	May	June	July	Aug.	Sep.	Oct.
<i>Vaccinium virgatum</i>	Blueberry		xx	xx						
<i>Cucumis sativus</i>	Cucumber				xx	xx	xx			
<i>Cosmos bipinnatus</i>	Cosmos Sensation			xx	xx	xx	xx	xx	xx	xx
<i>Coreopsis lanceolata</i>	Lance Leaf Tickseed			xx	xx	xx	xx	xx	xx	xx
<i>Echinacea purpurea</i>	Purple Coneflower					xx	xx	xx	xx	
<i>Fagopyrum esculentum</i>	Buckwheat			xx	xx	xx	xx	xx	xx	
<i>Gaillardia pulchella</i>	Indian Blanket Flower				xx	xx	xx	xx	xx	xx
<i>Lobularia maritima</i>	Sweet Alyssum		xx	xx	xx	xx				
<i>Lupinus perennis</i>	Wild Perennial Lupine			xx	xx	xx				
<i>Monarda punctata</i>	Spotted Beebalm				xx	xx	xx	xx	xx	xx
<i>Nemophila menziesii</i>	Baby Blue Eyes		xx	xx	xx	xx				
<i>Rudbeckia hirta</i>	Black-Eye Susan					xx	xx	xx	xx	xx
<i>Trifolium incarnatum</i>	Crimson Clover		xx	xx	xx	xx				