

Soil Biology Report Performed By:

Lab name: Sun Sprout Farm
 437 40th Ave
 Dunnell, MN, 56127
 Email: baileybeth21@gmail.com
 Phone: 8085576166
 Website:



Client:

Name: Bailey Scott
 Organization: Sun Sprout Farm
 437 40th Ave
 Dunnell MN 56127
 Email: baileybeth21@gmail.com
 Date Observed: 05-11-2025

Sample Name: Dorr's
Sample Type: Soil
Plants Present/Desired: soybeans
Plant Succession: Productive Pastures, Row Crops

Beneficial Microorganisms

	Recommended Range	Sample Results	
Fungi (ug/g)	135 - 1,350	11	Low: The fungal biomass is below the recommended minimum level for your plant's stage in succession. Please contact your Soil Biology Consultant.
Standard Deviation		16	Few target organism were present and variability was very high. Precision is very low.
Bacteria (ug/g)	135 - 1,350	324	Good: The bacterial biomass is within the recommended range for your plant's stage in succession.
Standard Deviation		50	Distribution of the target organisms in the sample was uniform; variation was small.
Actinobacteria (ug/g)	1 - 4	0.56	Low: The actinobacterial biomass is below the expected range. This is not a problem.
Standard Deviation		0.78	Few target organism were present and variability was very high. Precision is very low.
F:B Ratio	0.9:1 - 2:1	0.03	The F:B ratio is low. Increase fungal biomass or reduce bacterial biomass, and check predators to assess balance. Please contact your Soil Biology Consultant.

Minimum Value

Protozoa (Total)	> 50,000	11,413	Low: The number of beneficial protozoa is below the minimum requirement. Please contact your Soil Biology Consultant.
Standard Deviation		25,520	Few target organism were present and variability was very high. Precision is very low.
Flagellate (#/g)	(See Total)	11,413	
Standard Deviation		25,520	
Amoebae (#/g)	(See Total)	0	
Standard Deviation		0	

Nematodes

Bacterial-feeding (#/g)	300	0	None detected: Bacterial-feeding nematodes help keep bacterial populations in balance and enhance nutrient cycling.
Fungal-feeding (#/g)	100	0	None detected: Fungal-feeding nematodes help to release nutrients from fungal hyphae to the plants.
Predatory (#/g)	100	0	None detected: Predatory nematodes help reduce root-feeding nematode numbers.

Detrimental Microorganisms

Disease-Causing Fungi	Maximum Value	Sample Results	
Oomycetes (ug/g)	0	0	None detected: No disease-causing fungi were observed in the sample. Great!
Standard Deviation		0	Distribution of the target organisms in the sample was uniform; variation was small.

Anaerobic Protozoa

Ciliate (#/g)	0	0	None detected: No ciliates were observed in the sample. Aerobic conditions prevail. Great!
Standard Deviation		0	Distribution of the target organisms in the sample was uniform; variation was small.

Nematode

Root-feeding (#/g)	0	0	None detected: No root-feeding nematodes were observed. Great!
--------------------	---	---	--

Additional Comments: