

**Soil Biology Report Performed By:**

Lab name:  
 , ,  
 Email:  
 Phone:  
 Website:

**Client:**

Name: Bailey Scott  
 Organization: Sun Sprout Farm  
 437 40th Ave  
 Dunnell MN 56127  
 Email: baileybeth21@gmail.com  
 Date Observed: 05-09-2025

**Sample Name: organic plot n. crick**  
**Sample Type: Soil**  
**Plants Present/Desired: corn**  
**Plant Succession: Productive Pastures, Row Crops**

**Beneficial Microorganisms**

	Recommended Range	Sample Results	
Fungi (ug/g)	135    1,350	34	Low: The fungal biomass is below the recommended minimum level for your plant's stage in succession. Please contact your Soil Biology Consultant.
Standard Deviation		58	Few target organism were present and variability was very high. Precision is very low.
Bacteria (ug/g)	135    1,350	1,091	Good: The bacterial biomass is within the recommended range for your plant's stage in succession.
Standard Deviation		80	Distribution of the target organisms in the sample was uniform; variation was small.
Actinobacteria (ug/g)	1        4	0	Low: The actinobacterial biomass is below the expected range. This is not a problem.
Standard Deviation		0	Distribution of the target organisms in the sample was uniform; variation was small.
F:B Ratio	0.9:1    2:1	0.03	The F:B ratio is low. Increase fungal biomass or reduce bacterial biomass, and check predators to assess balance. Please contact your Soil Biology Consultant.

	Minimum Value		
Protozoa (Total)	> 50,000	34,238	Low: The number of beneficial protozoa is below the minimum requirement. Please contact your Soil Biology Consultant.
Standard Deviation		51,040	Few target organism were present and variability was very high. Precision is very low.
Flagellate (#/g)	(See Total)	34,238	
Standard Deviation		51,040	
Amoebae (#/g)	(See Total)	0	
Standard Deviation		0	

**Nematodes**

Bacterial-feeding (#/g)	300	0	None detected: Bacterial-feeding nematodes help keep bacterial populations in balance and enhance nutrient cycling.
Fungal-feeding (#/g)	100	0	None detected: Fungal-feeding nematodes help to release nutrients from fungal hyphae to the plants.
Predatory (#/g)	100	0	None detected: Predatory nematodes help reduce root-feeding nematode numbers.

## Detrimental Microorganisms

Disease-Causing Fungi      Maximum Value      Sample Results

Oomycetes (ug/g)	0	0	None detected: No disease-causing fungi were observed in the sample. Great!
Standard Deviation		0	Distribution of the target organisms in the sample was uniform; variation was small.

Anaerobic Protozoa

Ciliate (#/g)	0	0	None detected: No ciliates were observed in the sample. Aerobic conditions prevail. Great!
Standard Deviation		0	Distribution of the target organisms in the sample was uniform; variation was small.

Nematode

Root-feeding (#/g)	0	0	None detected: No root-feeding nematodes were observed. Great!
--------------------	---	---	--

**Additional Comments:**