

Soil Biology Report Performed By:

Lab name:

Email:
Phone:
Website:

Client:

Name: Bailey Scott
Organization: Sun Sprout Farm
437 40th Ave
Dunnell MN 56127
Email: baileybeth21@gmail.com
Date Observed: 07-26-2025

Sample Name: organic corn
Sample Type: Soil
Plants Present/Desired: corn
Plant Succession: Productive Pastures, Row Crops

Beneficial Microorganisms

	Recommended Range		Sample Results	
Fungi (ug/g)	135	1,350	39	Low: The fungal biomass is below the recommended minimum level for your plant's stage in succession. Please contact your Soil Biology Consultant.
Standard Deviation			87	Few target organism were present and variability was very high. Precision is very low.
Bacteria (ug/g)	135	1,350	300	Good: The bacterial biomass is within the recommended range for your plant's stage in succession.
Standard Deviation			17	Distribution of the target organisms in the sample was uniform; variation was small.
Actinobacteria (ug/g)	1	4	0.6	Low: The actinobacterial biomass is below the expected range. This is not a problem.
Standard Deviation			0.6	Few target organism were present and variability was very high. Precision is very low.
F:B Ratio	0.9:1	2:1	0.13	The F:B ratio is low. Increase fungal biomass or reduce bacterial biomass, and check predators to assess balance. Please contact your Soil Biology Consultant.

	Minimum Value		
Protozoa (Total)	> 50,000	45,651	Low: The number of beneficial protozoa is below the minimum requirement. Please contact your Soil Biology Consultant.
Standard Deviation		47,743	Few target organism were present and variability was very high. Precision is very low.
Flagellate (#/g)	(See Total)	45,651	
Standard Deviation		47,743	
Amoebae (#/g)	(See Total)	0	
Standard Deviation		0	

Nematodes			
Bacterial-feeding (#/g)	300	0	None detected: Bacterial-feeding nematodes help keep bacterial populations in balance and enhance nutrient cycling.
Fungal-feeding (#/g)	100	0	None detected: Fungal-feeding nematodes help to release nutrients from fungal hyphae to the plants.
Predatory (#/g)	100	0	None detected: Predatory nematodes help reduce root-feeding nematode numbers.

Detrimental Microorganisms

Disease-Causing Fungi	Maximum Value	Sample Results	
Oomycetes (ug/g)	0	0	None detected: No disease-causing fungi were observed in the sample. Great!
Standard Deviation		0	Distribution of the target organisms in the sample was uniform; variation was small.

Anaerobic Protozoa

Ciliate (#/g)	0	0	None detected: No ciliates were observed in the sample. Aerobic conditions prevail. Great!
Standard Deviation		0	Distribution of the target organisms in the sample was uniform; variation was small.

Nematode

Root-feeding (#/g)	0	0	None detected: No root-feeding nematodes were observed. Great!
--------------------	---	---	--

Additional Comments: