


Pineapple News

Issue No. 17 Newsletter of the Pineapple Working Group, International Society for Horticultural Science May, 2010

Table of Contents

| | |
|--|-----------|
| News of General Interest..... | 2 |
| Maui Pineapple Company Closes..... | 2 |
| 'MD-2' Pineapple Named the American Society for Horticulture Science 2010 Outstanding Fruit Cultivar..... | 2 |
| 7 th International Pineapple Symposium 2010..... | 3 |
| Pineapple Working Group..... | 4 |
| Contribute to Pineapple News..... | 4 |
| News From Belgium..... | 4 |
| Pre and Postharvest Metabolism of Leaves on the Pineapple Fruit Crown | 4 |
| News From Brazil..... | 8 |
| Alternative Control of Pineapple Fusariosis on Irrigated 'MD-2' Pineapple in Brazil..... | 8 |
| News From Cuba..... | 9 |
| Chemical Sterilization for Propagation of Pineapple Plantlets in Temporary Immersion Bioreactor..... | 9 |
| News From France..... | 15 |
| Growth Indicators for Queen Victoria Pineapple Versus Sum of Temperatures, Basis for a Heat-unit Model of Vegetative Growth..... | 15 |
| A Bait and Trap Method for Sampling Symphylid Populations in Pineapple..... | 18 |
| News from Ghana..... | 23 |
| Pineapple and Carbon Emissions..... | 23 |
| News from Malaysia..... | 23 |
| Handling of Fresh-cut Pineapple for Fresh Consumption..... | 23 |
| News From Taiwan..... | 24 |
| RAPD Marker Assisted Selection of EMS Induced Pineapple (<i>Ananas comosus</i> (L.) Merr.) Mutants..... | 24 |
| News From the United States..... | 27 |
| Role of Proteases During Ripening of Pineapple Fruit..... | 27 |
| Environmental Friendly Approaches for Managing Nematodes and Weeds on Pineapple..... | 27 |
| Commonalities of Pineapple, Agaves, and Cacti..... | 32 |
| Services..... | 33 |
| Commercial Services..... | 33 |
| Professional Services..... | 34 |
| Book Reviews and Web Sites..... | 34 |
| Book Reviews..... | 34 |
| Web Sites of Possible Interest..... | 34 |
| New References on Pineapple..... | 35 |
| Instructions to Contributors to Pineapple News..... | 41 |



News of General Interest, Pineapple Working Group News

Dear Colleagues:

The big news this year is the 7th International Pineapple Symposium, which is being held in Johor, Malaysia in July 2010. As of April 19, 2010, the Secretariat of the symposium reported they had received 32 oral and 105 poster papers. I am looking forward to attending this year's symposium and hope to meet many of you there.

Maui Pineapple Company Closes

Many of us associated with the Hawai'i pineapple industry were saddened to read on Tuesday, November 3, 2009 that Maui Pineapple Company planned to cease planting of pineapple immediately and would shut down all operations at the end of the year (http://www.hawaiimagazine.com/blogs/hawaii_today/2009/11/4/Maui_Land_Pineapple_production_ends). The last harvest was to be made on December 23, 2009. Not long after the closure was announced, a news article reported that a group of investors planned to continue growing fresh pineapples on Maui under the name Haliimaile Pineapple Company, Ltd. The new company is to lease land from Maui Land and Pineapple Company. The organizers of the new company included five former Maui Pineapple Company executives and former Maui Pineapple Company manager Darren Strand will take over as president of the new company. The news release (<http://www.mauinews.com/page/content.detail/id/527256.html>) indicated that the new company would concentrate on the island market with a small percentage of the fruit to be shipped to premium markets on the U.S. mainland. The company plans to grow Maui Gold® pineapple on about 1,000 acres in the Haliimaile area of Maui. The company expected to save about 65 agricultural jobs with the hope that more could be added in the future.

As of this writing, Dole continues to grow pineapple on about 3,000 acres on the island of Oahu, which is where the main market is located. Dole also ships a limited amount of pineapple to California and other west-coast markets.

'MD-2' Pineapple Named the American Society for Horticulture Science 2010 Outstanding Fruit Cultivar

On reading an announcement in 2009 about the American Society for Horticultural Science Outstanding Fruit Cultivar Award, Duane Bartholomew and Robert Paull spearheaded an effort to nominate 'MD-2' for this award. The nomination reads in part: The cultivar MD-2 resulted from a cross made in 1970 by Dr. David D. F. Williams, plant breeder and director of the Pineapple Research Institute of Hawaii (PRI), a nonprofit research institute supported by the Hawaii Pineapple Growers Association (HPGA). PRI was funded at the time the cross was made by Del Monte Hawaii Inc., Dole Food Co., Hawaiian Fruit Packers, Libby McNeil & Libby, Maui Pineapple Co. and several smaller pineapple growers. Dr. Williams' able assistants in the pineapple breeding program were Frank Bermudas and Toshio Minagawa. The nomination called attention to the outstanding contributions to pineapple breeding made by the Pineapple Research Institute of Hawaii and to David D.F. Williams. The justification for naming 'MD-2' as the outstanding fruit cultivar in 2010 was based on the cultivars overwhelming success in the international pineapple markets.

Dr. Williams, now in his early 80s, and his daughter Wendy flew to St. Louis, Missouri to accept the award. Photos of the award ceremony did not survive a camera accident so the photo in Figure 1 was taken at a friends home in the beautiful Rocky Mountains in Colorado. Dr. Williams has been retired for more than 10 years and he and his wife Elsa currently reside near a number of their children in Fort Collins, Colorado, U.S.A.




Figure 1. Photo of Dr. David D.F. Williams holding the medal recognizing 'MD-2' pineapple as the American Society for Horticulture Science 2010 Outstanding Fruit Cultivar. The medal was awarded to Dr. Williams and the inscription reads: 'MD-2' Pineapple. Dr. David D.F. Williams and Forbears, Pineapple Research Institute of Hawaii.

7th International Pineapple Symposium 2010



7TH INTERNATIONAL PINEAPPLE SYMPOSIUM 2010



Malaysia Hosts the 7th International Pineapple Symposium (7th IPS)

The Symposium Organizing Committee, Malaysia Agriculture Research and Development Institute (MARDI), Malaysian Pineapple Industry Board (MPIB) and International Society of Horticultural Science (ISHS), invites all of you to the 7th IPS to be held from 13 to 15 July 2010. The theme "GLOBAL PINEAPPLE INDUSTRY- THE WAY FORWARD" emphasizes on the latest trends and development in the global pineapple sector. All aspects relating to pineapple cultivation (industry and trade, breeding, genetics, plant physiology and cultural practices, pest and disease management, post harvest handling and product development, consumer and marketing) will be covered, taking into account the most recent scientific and technical advancement around the world.

The symposium will be held at PERSADA Johor International Convention Center in Johor Baru, Malaysia. The state of Johor is known for its pineapple cultivation on peat soil since 1960s. Being tropical and evergreen, Johor Baru with its beautiful landscape, just across the causeway from Singapore is an excellent meeting place for scientists, professional, industry players, traders, students and any one that has interest in pineapple business.

About 200 participants locally and from all over the world and international organizations will participate in the 7th IPS 2010. Interesting abstracts are being submitted by participants locally and abroad covering various disciplines.

The scientific program which encompasses oral and poster presentations, exhibition and field visit, will provide delegates with information, a forum for discussion and networking opportunities for building beneficial relations with the dynamic pineapple industry.

Important dates for 7th IPS:

- Full paper submission- before or on 13th July 2010
- Paper presenters must pay their registration fees 2 months before the Symposium i.e. 13 May 2010.

We are looking forward to meeting you at the 7th IPS
Convener,
7th IPS

Jointly organized by:



- Malaysia Agriculture Research and Development Institute (MARDI)
- Malaysian Pineapple Industry Board (MPIB)
- International Society of Horticultural Science (ISHS)

In co-operation with:



- Department of Agriculture, Malaysia (DOA)
- Federal Agricultural Marketing Authority (FAMA)
- International Tropical Fruits Network (TFNet)

Supported by:



- Ministry of Agriculture and Agro-based Industry Malaysia (MOA)
- Tourism Malaysia

For further details please visit our website at
<http://www.mardi.my>

For the most current information on the symposium, please point your web browser to the symposium web site at <https://anjungnet.mardi.gov.my/Conference.nsf/PineApple?OpenPage>.

Pineapple Working Group of the International Society for Horticulture Science

I would like to remind readers of Pineapple News that all proceedings of the six international pineapple symposia are available for sale from the ISHS web site. The ISHS also provides free web access to the abstracts of all papers presented at the six symposia as well as to the abstracts of papers in all volumes of *Acta Horticulturae* published by the ISHS. There are many benefits to being a member of the ISHS and one that I have enjoyed is the opportunity to download at no additional cost a limited number of copies of articles published in *Acta Horticulturae*. The ISHS is one of the foremost organizations promoting cooperation and communication among horticultural researchers, growers and consumers. The ISHS continues to expand its offerings to members as well as to provide the structure under which our Pineapple Working Group (<http://www.ishs.org/science/T07.php>) functions. Detailed information about ISHS and the benefits of membership can be found at <http://www.ishs.org> or you can write to the ISHS Secretariat, P.O. Box 500, 3001 Leuven, Belgium (E-Mail: info@ishs.org).

Contribute to Pineapple News

Information on how to contribute to Pineapple News can be found at the end of the newsletter. You can also contact Duane Bartholomew, the editor, at duaneb@hawaii.edu. ♦

News From Belgium

Pre and Postharvest Metabolism of Leaves on the Pineapple Fruit Crown

Londers E.¹, Ceusters*J.¹, Godts C.¹, Van De Poel B.², De Proft M.P.¹

¹Division of Crop Biotechnics, Department of Biosystems, Faculty of Bioscience Engineering, Katholieke Universiteit Leuven, Willem De Croylaan 42, B-3001 Leuven, Belgium

²BIOSYST-MeBioS, Department of Biosystems, Faculty of Bioscience Engineering, Katholieke Universiteit Leuven, Willem de Croylaan 42, B-3001 Leuven, Belgium

*Corresponding author E-mail: johan.ceusters@biw.kuleuven.be; Phone, +32 16322403; Fax., +32 16322966.

Abstract

Crown burn on exported pineapple fruit causes serious economical losses. In this study, crown burn occurrence was hypothesized to be related to the Crassulacean Acid Metabolism (CAM). However, the functioning of this metabolism in the crown leaves remained unexplored. Therefore, pre and post harvest bio-activity of the crown leaves of pineapple fruit were studied. With the fruit still attached to the plant, CAM activity was detected in the crown leaves. While post-transport diurnal light cycle conditions seemed to reactivate CAM in the crown leaves, no CAM activation was registered during cold-storage transport under dark conditions (simulating transport). The substantial organic acid increase in the crown leaves observed during transport is thought to originate from the pineapple fruit itself.

Introduction

As a part of the grading standard, not only pineapple fruit but also crown quality is an aesthetic characteristic of economic concern. As such, leaf damage, occurring as brown spots on the crown leaves, is causing economical losses. The authors hypothesized these spots to be the result of physiological disorders in a similar way as earlier described for the ornamental bromeliad *Aechmea*. In these ornamental plants the build-up of high organic acid concentrations causes chlorenchyma cells to rupture (Londers et al., 2005b; Londers, 2006).

Nocturnal organic acid build-up is a central feature of CAM (Crassulacean Acid Metabolism) with pineapple being the most important crop exhibiting this metabolism. Stomata in mature leaves open at night and CO₂ is assimilated into malate by the carboxylating enzyme phosphoenolpyruvatecarboxylase (PEPC). During daytime CO₂ will be released and processed via Rubisco behind closed stomata. The pineapple crown is a continuation of the vegetative stem, and the spirally arranged leaflets have a similar morphology. However, the photosynthetic activity of crown leaves remained unexplored.

Therefore the authors investigated the metabolism of the crown leaves both in the field and under post harvest transport conditions. To figure out on plant carbon metabolism, nocturnal organic acid build-up is often used as a key determinant for CAM (Winter and Smith, 1996). Furthermore, the stomata opening index gives a good indication for gas exchange patterns (Londers et al., 2005b), with nocturnal gas exchange characteristic for CAM. These measurements can further be refined by gas exchange data about leaf CO₂ exchange. The results are presented and discussed in the broader context of crown leaf damage.

Materials and methods

Experimental setup and plant material

The experimental setup is summarized in Table 1. The metabolism of the pineapple crown leaves was studied at different stages in the pineapple production chain. For the preharvest stage, the metabolism was studied by determining the stomata opening index of the leaves of the pineapple crown. Therefore, *Ananas comosus* hybrid plants were grown in the greenhouse of the Katholieke Universiteit Leuven under greenhouse conditions as described in Londers et al. (2005b). For the postharvest transport analyses, fruits (*Ananas comosus* hybrid) were obtained from a commercial pineapple plantation in Ecuador. Pineapple crown leaves were sampled at harvest and at their arrival in the port of Antwerp, Belgium, after a transport period of three weeks in dark and cooled conditions (about 8 °C). Finally, storage simulation experiments were executed, by placing fruits in cooled (10 °C) climate chambers under both a continuous dark and a diurnal light cycle (12/12) regime. Under the diurnal light cycle regime, light intensity in the climate chamber averaged around 150 mol m⁻² s⁻¹.

Stomata opening index

The stomata index was the percentage of opened stomata of hundred stomata examined. Fully developed crown leaves were sampled on the abaxial leaf side between the half leaf length and the leaf tip. After removing trichomes with adhesive tape, a thin coat of transparent nail polish was layered onto the leaf surface and allowed to dry for 15 min. Dry polish coats were carefully removed using transparent adhesive tape (Scotch crystal clear) and stuck on glass slides for microscopic observations (Londers, 2006).

Organic acid measurements

Samples for organic acid analysis were collected from fully developed leaves of the pineapple crown (n = 8 fruits). Samples were subsequently weighed, immediately frozen in liquid nitrogen to arrest enzyme activity and stored at -20 °C until required for analysis. The samples were extracted with water after grinding nitrogen frozen tissues with a tissue grinder. For the determination of total titratable acids, extracts were titrated (Metrohm, 702 SM Titrino) to an end point of pH 8.1 using 0.01 N KOH. Extraction and quantification of malic acid was accomplished as described by Londers et al. (2005b) using high performance liquid chromatography (Waters 510, Waters, Milford, MA, USA) with detection at 210 nm (Waters 484, Waters, Milford, MA, USA) using an aminex HPX87-H (300 mm x 7.8 mm) resin-based column from Bio-Rad (Hercules, CA, USA).

Table 1. Experimental setup and actions

| Sampling points in production chain | Time of sampling (days) | Climate conditions | Type of measurement |
|-------------------------------------|-------------------------|----------------------------|----------------------------|
| Field | Before harvest | | Stomata opening index |
| Harvest | 0 | | Total titratable acids |
| Transport | 21 | Dark + 8°C | Total titratable acids |
| Storage | 28 | Dark + 10°C | Total titratable acids |
| | | | Net CO ₂ uptake |
| | | Diurnal light cycle + 10°C | Total titratable acids |
| | | | Net CO ₂ uptake |

Gas exchange measurements

Leaf CO₂ exchange data were collected on the youngest fully expanded leaf from the pineapple crown under a continuous dark and alternated light/dark (12/12) regime. A LCi Portable Photosynthesis System (ADC BioScientific Ltd., UK) was used. The top part of the leaf was enclosed in the leaf chamber. Short-term fluctuations in the CO₂ concentration of the incoming air were buffered by passing the air through a 25 L metal bottle. Gas exchange data were collected using a 10 min interval (n = 5 fruits).

Where appropriate, the data were analyzed using the statistical software package SAS Enterprise Guide 4.0. Before carrying out statistical tests normality of the data was checked by means of the Kolmogorov-Smirnoff statistic (p > 0.05). Means were compared by two sample t-test or by Tukey's studentized range test.

Results

Preharvest metabolism of leaves

The pattern of stomatal opening of the crown leaves (Figure 1) with the fruit still attached to the plant was typical of CAM plants. Stomata of CAM plants remain mainly closed during daytime and open at night, controlled by internal CO₂ concentrations (Salisbury and Ross, 1991).




Figure 1. Stomata opening index (% open stomata) of pineapple crown leaves sampled before harvest. The curve is representative of 10 measurements with SE < 15 %.

Postharvest metabolism of leaves

Total titratable acids in the crown leaves at some critical points in the pineapple fruit production chain, starting from the time of harvest, can be seen in Table 2. Relative to leaves sampled at harvest, total titratable acid values almost doubled during dark, cold transport. The post-transport treatments evaluated did not affect the organic acid levels. There was a significant accumulation of malic acid in pineapple crown leaves after 7 days of post-transport treatment when crowns were exposed to a diurnal light cycle cold-regime (Figure 2). Under dark cold conditions, CAM seems not to be active. Net CO₂ uptake under different post-transport treatment conditions (Figure 3) supports this statement. While there is a clear pattern of CO₂ uptake during the night in a 12/12 light-dark diurnal cycle, it is absent under continuous dark conditions.

Table 2. Total titratable acids (Eq gfw⁻¹) in the crown leaves of pineapple fruit at critical points in the production chain.

| Time of sampling (days after harvest) | Treatment [†] | Leaf total titratable acids (Eq gfw ⁻¹) |
|---------------------------------------|------------------------|---|
| 0 | --- | 162 |
| 21 | OT | 308 ± 22 a |
| 28 | OT + Dark + 10°C | 299 ± 18 a |
| 28 | OT + DLC + 10°C | 288 ± 12 a |

[†]OT = Overseas Transport; DLC = Diurnal Light Cycle. Leaves were sampled at 8:00 AM. SE (n = 8 fruits) and Tukey's Studentized Range Test (P < 0.05). Treatments followed by the same letter were not significantly different from each other.

Discussion and conclusions

Crown burn during overseas transport of pineapple fruit is a problem of economic concern. Visually, crown burn seems to be identical to a specific type of leaf damage on the leaves of many ornamental *Aechmea* cultivars, which was studied earlier by the authors (Londers et al., 2005a). Microscopic analysis of damaged crown leaves (results not shown) revealed ruptured cells in the chlorenchyma of the leaves, which matches damage images obtained earlier for *Aechmea* (Londers et al., 2005a). Based on both the visual and microscopic images, the authors hypothesized crown burn to be the result of physiological disorders similar to those described earlier for *Aechmea* (Londers et al., 2005b; Londers et al., 2006; De Proft et al., 2007). More specifically, high organic acid concentrations in the leaves due to CAM might act as water pumps, resulting in (lethal) leaf turgor pressure. Especially *Aechmea* plants kept under prolonged dark conditions seem to be at high risk for this type of leaf damage, due to a disturbed CAM pattern (Ceusters, 2008). To evaluate the mechanism described above for the crown burn case, the metabolism of the crown leaves and especially leaf organic acid concentrations during overseas transport needed to be investigated.

With the pineapple crown being a continuation of the vegetative stem leaves, and the spirally arranged leaflets having a similar morphology, the authors suppose the crown to act as CAM. This was supported by diurnal stomata opening analyses (Figure 1) with the fruit still attached to the plant, which clearly indicated CAM.




Figure 2. Nocturnal malic acid build-up in pineapple crown leaves from 5:00 p.m. to 8:00 a.m. after 7 days of post-transport treatment (n = 8 fruits). Diurnal light cycle at 10 °C (black bars); continuous dark at 10 °C (white bars). Statistically significant nocturnal malic acid build-up (*P<0.05) is indicated by *.

Concerning the organic acid concentrations in the crown leaves, total titrable acids doubled during the three weeks cold-transport period (Table 2). Nanayakkara et al. (2005) already described the increase of total titrable acids in the crown leaves during cold-storage.

Though, storage under diurnal light cycle conditions seems to activate a diurnal stomatal rhythm with a similar pattern of CO₂ gas exchange (Figure 2) and nocturnal accumulation of malate in the leaves (Figure 3).

To conclude, the increase of total titrable acids in the leaves on the crown of pineapple fruit during dark cold-transport is not believed to be caused by a disturbed CAM. Total titrable acids seem to build-up gradually during the overseas cold-transport period (results not shown). Possibly this increase in total titrable acids in the crown originates in the migration of organic acids from the fruit into the crown. Therefore, CAM interacting measures will not affect the risk of crown burn. Though, one should be aware that the availability of external water should be absolutely avoided throughout the whole postharvest chain. External water restriction limits the risk of lethal turgor pressure build-up, driven by organic acid concentrations in the leaves which are increasing during transport.

Funding




Figure 3. Postharvest net CO₂ uptake (µmol m⁻² s⁻¹) pattern for young fully expanded pineapple crown leaves under a continuous dark (■) and diurnal light cycle (12/12) (□) cold-regime. Both curves are representative of five replicate runs with SE < 10 %.

This research was supported by the Institute for the Promotion of Innovation through Science and Technology in Flanders (IWT-Vlaanderen).

References

- Ceusters, J., 2008. Plasticity of CAM in *Aechmea*: possibilities and limitations. Doctoral thesis, Katholieke Universiteit Leuven, Leuven.
- De Proft, M.P., Ceusters, J., Londers, E., 2007. Leaf quality management of *Aechmea* throughout the supply chain. *Acta Hort.* 755, 39-44.
- Londers, E., Vervaeke, I., Deroose, R. and De Proft, M.P., 2005a. Leaf Quality of *Aechmea* under Non-Adapted Greenhouse Conditions. *Acta Hort.* 669, 275-282.
- Londers, E., Ceusters, J., Vervaeke, I., Deroose, R., De Proft, M.P., 2005b. Organic acid analysis and plant water status of two *Aechmea* cultivars grown under greenhouse conditions: implications on leaf quality. *Scientia Hort.* 105, 249-262.
- Londers, E., 2006. Leaf damage in CAM bromeliads: a hydrophysiological approach. Doctoral thesis, Katholieke Universiteit Leuven, Leuven.
- Nanayakkara, K.P.G.A., Herath, H.M.W. en Senanayake, Y.D.A., 2005. Effects of Post-harvest Treatments of Potassium, Malic Hydrazide, Paraffin-Polyethylene Wax and Light on Reducing Internal Browning (IB) and Malic Acid Content in Pineapple (*Ananas comosus* (L.) Merr. Cv Mauritius) during Cold-storage. *Acta Hort.* 666, 329-335.
- Salisbury, F.B., Ross, C.W., 1991. *Plant Physiology*. Wadsworth Publishing Company, California, USA.
- Winter, K., Smith, J.A.C., 1996. An Introduction to Crassulacean Acid Metabolism. *Biochemical Principles and Ecological Diversity*. In: Winter, K., Smith, J.A.C. (Eds.), *Crassulacean Acid Metabolism : Biochemistry, Ecophysiology and Evolution*. Springer-Verlag, Berlin-Heidelberg, Germany, New York, USA. pp. 1-10.◆

News From Brazil

Alternative Control of Pineapple Fusariosis on Irrigated 'MD-2' Pineapple in Brazil

Remulo A. Carvalho¹, Eliazar F. de Oliveira¹, José T. de Lacerda¹, Miguel B. Neto¹, Paulo de T. de M. Duarte² and Rosimar de Figueiredo²

¹ Agronomists - EMEPA (Paraíba State Agricultural Research Company)

² Agronomists - AGRIVALE (Pineapple Producing and Exporting Company)

EMEPA-PB, Rua Eurípedes Tavares, 210, CEP: 58.013-290, João Pessoa, PB, Brazil

Email: remuloc@hotmail.com

Brazil is the biggest pineapple (*Ananas comosus* var. *comosus*) producer in the world with high yields and excellent fruit quality. However, despite the development of technologies for the pineapple crop, huge losses still occur due to high incidences of fusariosis disease, a devastating fruit rot caused by the fungus *Fusarium subglutinans*, which attacks not only the fruit, but also the whole plant and its slips which are used as propagating material in planting. In Brazil, 'Pérola' and 'Smooth Cayenne' plants are susceptible to fusariosis and 'MD-2' is extremely susceptible and the disease is difficult to control in irrigated commercial fields. Traditionally, the control of this disease is based on preventive applications of chemical fungicides. Weekly applications of a mixture of benzimidazole fungicides and carbamate insecticides (to control a fruit pest which occasionally opens the way for the fungus) during the 4 to 5 week period in which the pineapple flowers are open efficiently control fusariosis. Usually excessive residues of pesticides make exportation of the fruit impossible. On the other hand, there is increasing concern about the possible impact of fungicides on the environment and the onset of resistance to chemical crop protection agents together with a consumer preference for chemical-free produce, which has led to a search for agriculture products treated with natural substances. Therefore, the objective of this research was to identify an alternative control of pineapple fusariosis. Different dosages of citric extracts, food preservatives, pyroligneous acid and tannins of cultivated *Acacia mearnsii* were evaluated. The tannins are extracted and commercialized worldwide for utilization on the leather and tanning industries. Four experiments involving almost 15,000 plants were carried out in a private pineapple farm located in the semi-arid zone of the state of Bahia, Brazil in 2008. The experimental design was of randomized blocks with 8 treatments and four replicates. The tannins were applied once and twice a week at 5, 6, and 7% dosages. Most of the alternative products were inefficient against this disease except for the tannins of *A. mearnsii*. Although there was no difference among tannins treatments the difference in relation to the control was very impressive considering the favorable environment to the disease. The fruits were evaluated immediately after the harvest in search of symptoms of fusariosis rot. In this experiment the incidence of fusariosis was reduced from 72% in the control treatment (Figure 1A) to 22% in some tannin treatments (Figure 1B).



Figure 1. Fruits from untreated control infected with *Fusarium subglutinans* (A) and fruits with fusariosis from a comparable plot treated with black wattle tannins (B).



News From Cuba

Chemical Sterilization for Propagation of Pineapple Plantlets in Temporary Immersion Bioreactor

M. Escalona¹, Z. Fundora¹, C. Aragón¹, I. Capote¹, N. Castañedo², D. Pina¹ and J. González-Olmedo¹.

¹ Laboratory for Plant Cell and Tissue Culture, Bioplant Centre, University of Ciego de Avila, 69450, Cuba. E-mail: mescalona@bioplantac.cu; Fax: 53-33 266340.

² Centro de Bioactivos Químicos, Universidad Central, Villa Clara, Cuba

Summary

Vitrofur G-1 (patent no. 22676) is a relatively new, broad spectrum biocide derived from sugarcane plants, used in the medical industry and in plant tissue culture. The active ingredient is 1-(5 bromofur-2-IL)-2-bromo-a-nitroeteno. G-1 is effective against both bacteria and fungi; it is heat stable and can be used in the culture medium without any toxic effect on plant tissue. These characteristics of G-1 make it an attractive alternative for the control of contaminations during temporary immersion culture. In this context, the main objective of this study was to establish a procedure for pineapple propagation in temporary immersion bioreactors using G-1, as way of medium chemical sterilization, as well as to evaluate net photosynthesis and quality of shoots prior to acclimatization (elongation phase) using this procedure.

Key words: pineapple, Temporary Immersion, Bioreactor, photosynthesis, liquid medium.

Introduction

In order to overcome the current limitations of micropropagation, different procedures have been developed. Among them, the temporary immersion of explants in a liquid medium has been achieved by using different bioreactors, which are described and grouped into four categories according to operation. All of these systems have as advantages the avoidance of continuous immersion, which adversely affects growth and morphogenesis, provision of adequate oxygen transfer, sufficient mixing, and limited shear levels. The technique enables sequential medium changes and automation while reducing contamination and costs (Etienne and Berthouly 2002).

In vitro culture methods of pineapple plants allows for a more rapid increase of selected cultivars than traditional propagation methods. Commercial pineapple micropropagation involves sequential culturing in liquid medium for meristem and axillary shoot bud multiplication (Daquinta and Benegas, 1997). Using this approach, annual pineapple production is limited as a result of the number of pineapple plants needed annually to start up new plantations. In general, the commercial use of micropropagation is currently reduced because of high production costs. It is due primarily from high labour cost, low multiplication rate, and poor survival rates during acclimatization.

A procedure for the mass propagation of pineapple plants (*Ananas comosus* L. Merr cv Smooth cayenne) using a temporary immersion technique was described by Escalona *et al.*, (1999). This procedure involved three distinct phases in the semi-automated temporary immersion system: shooting, bud differentiation and elongation. An efficient and cost-effective method for commercial

micropropagation of 'Smooth Cayenne' pineapple using a periodic immersion bioreactor was also developed by Firoozabady and Gutterson (2003). Both methods are applicable to the micropropagation of standard cultivars but are also applicable to the rapid scale-up of planting material of a newly developed cultivar.

Microbial and fungal contamination is one of the most persistent problems in plant cell and tissue culture. Surface removal of contaminating organisms from plant tissue being prepared for culture is generally by treatment with oxidizing hypochlorite solutions, ethanol, mercuric chloride or a combination of these or other treatments. Antibiotics are used, but are not always effective, can alter plant growth, are costly, and resistant strains can result with extensive use (Niedz 1998). Plant Preservative Mixture (PPM) has been routinely added to tissue culture medium to control air and waterborne bacterial and fungal contaminants effectively (Niedz 1998; George and Tripepi 2001; Beruto *et al.*, 2004).

Vitrofur G-1 (patent no. 22676) is a relatively new, broad spectrum biocide derived from sugarcane plants, which has been used in medical industry and plant tissue culture. The active ingredient is 1-(5-bromofur-2-yl)-2-bromo-4-nitroethene. G-1 is effective against both bacteria and fungi, is heat stable and can be used in the culture medium without any toxic effect on plant tissue (Alvarado *et al.* 1997). These characteristics of G-1 make it an attractive alternative for use in temporary immersion culture. In this sense, the main objective of this study was to establish a procedure for pineapple propagation in temporary immersion bioreactor (TIB) using G-1, as way of medium chemical sterilization, as well as to evaluate some physiological parameters of pineapple shoot elongation in TIB using this procedure.

Material and Methods

Plant Material

Pineapple plantlets (*Ananas comosus* L. cv Smooth Cayenne) were obtained from established liquid cultures grown on a shooting medium, which consisted of MS (Murashige & Skoog 1962) salts supplemented with 9.32 μ M benzyladenine (BA) and 1.61 μ M naphthaleneacetic acid (NAA), as recommended by Daquinta and Benega (1997). The cultures were grown under a 16-h photoperiod from cool-white fluorescent lamps (80 μ mol m⁻² s⁻¹), at 25 \pm 1.0 °C.

Culture conditions

The Temporary Immersion Bioreactor (TIB) used was described by Escalona *et al.* (1999). Five explants (*in vitro*-cultured plants with two small shoots) approximately 2 to 3 cm in length were cultured in each container. The degree of endogenous contamination in explants was determined by culturing each explant on nutrient broth medium for 72 h at 37 °C before inoculation in the TIB. The culture vessels used for the experiment were 10.5 cm in diameter, 20 cm high and had a volume of 1000 mL; 200 mL of liquid medium was added to the culture vessel for each explant (Escalona *et al.*, 1999). The basal medium consisted of MS salts with 30 g L⁻¹ sucrose and 1.0 mg L⁻¹ thiamine. The medium was sterilized using 28 mg L⁻¹ of 100% a.i. G-1, the amount recommended to reduce or eliminate airborne contamination in liquid media using the TIB (Fundora *et al.*, 2003). The control treatment was autoclaved at 121 °C for 30 min. The pH of the culture medium was adjusted to 5.8 after addition of G-1 and autoclaving, respectively. All cultures were kept in a growth room at a temperature of 24 °C under cool-white fluorescent lamps as above with a 16 h photoperiod.

Concentration of plant growth regulators in the medium for shoot proliferation

Different concentrations of the main plant growth regulators that are included in the proliferation medium for pineapple, established previously by Escalona *et al.*, (1999), were tested. A medium with 9.32 μ M BA, 1.61 μ M NAA, and 4.37 μ M PB was the control. BA levels assessed were 0, 2.33, 4.66, 6.99, and 9.32 μ M, all with the same composition of NAA and PB as described above. Once the best concentration of BA was determined, 4.37 μ M PB was added and NAA levels of 0, 0.54, 1.07 and 1.61 μ M NAA were tested. After the best concentrations of BA and NAA were determined, paclobutrazol (PB) [(2RS,3RS)-1-(4-chlorophenyl)-4,4-dimethyl-2-(1H-1,2,4-triazol-1-yl)pentan-3-ol] at 0.0, 1.09, 2.18, 3.28 and 4.37 μ M were tested. The experimental conditions were as established by Escalona *et al.*, (1999). Immersion frequency was 4 min every 3h. The duration of the proliferation phase was 30 d and 28 mg L⁻¹ of G-1 was also used to sterilize culture medium. Multiplication rate was determined by dividing the final number of shoots by the initial number of explants, and was evaluated after 30 d of culture.

Immersion frequency during proliferation and elongation phases

Based on the experiments described above, a proliferation medium consisting of MS formula with 6.66 μ M BA, 1.07 μ M NAA and 4.37 μ M PB was used. Then, different immersion frequencies of 1, 3 and 5 h were tested. For all treatments, the immersion time was 4 min. After a shoot-multiplication period (30 d), the medium was replaced with MS medium supplemented with 2.22 μ M BA and 4.06 μ M GA3. After seven days, the medium was replaced again with a MS medium supplemented with 4.06 μ M GA3 and shoots were grown for 21 d to promote shoot elongation. All of these steps were previously described by Escalona (1999), but in this experiment, were modified through the addition of 28 mg L⁻¹ G-1 to sterilize the culture medium. Multiplication rate, length of shoots, leaf area, fresh and dry mass and chlorophyll content were determined after the 21-day elongation phase. Leaf area was determined by a photogravimetric method (Sestak *et al.*, 1971). Chlorophyll a + b were extracted in acetone (80%) and chlorophyll contents were determined using the equations of Porras *et al.* (1991). Dry weight was measured after drying the shoots for 72 h at 70 °C.

Determination of photosynthetic parameters

For measurements of the maximum photosynthetic capacity (P_n), fully expanded leaves from shoots were used between 4-5 h after the beginning of the photoperiod. A maximum photosynthetic rate was measured using a portable CIRAS-2 photosynthesis System (Europe, PP Systems, UK). The whole area of the cuvette (PLC6) was covered with completely expanded young 2.5 cm² pineapple leaf. The carbon dioxide concentration and the humidity of the air entering the leaf chamber was 375 μmol mol⁻¹ and 80%, respectively, under a photosynthetic photon flux of 2000 μmol m⁻² s⁻¹. The measurements were done on five plants with 10 replicates for a total of 50 values. Prior to light measurements the maximal and stable photosynthesis for the pineapple plants was determined. At 600 μmol m⁻² s⁻¹, the photosynthesis was maximal and stable. Measurements were performed after 21d of culture prior to acclimatization (elongation phase).

Statistics

All statistical analyses were done using SPSS (Chicago, IL, USA, version 9.0). Differences due to hormone concentration, immersion frequency and shoot morphological parameters were analyzed using one-way ANOVA, followed by a Tukey comparison at p 0.05.

Results

Concentration of plant growth regulators in the medium for shoot proliferation

Three experiments were conducted to determinate the effect of different concentrations of the main plant growth regulators of pineapple in the presence of G-1 on multiplication rate and shoot quality. When different concentrations of BA were tested, the highest multiplication rate was achieved at 6.99 μM, which was not different from the autoclaved treatment. At the highest concentration of BA, shoot multiplication rate was decreased relative to the autoclaved treatment in the presence of G-1. Shoot multiplication rate in media with the best concentration of BA and without NAA, but with 4.37 μM PB increased significantly. However, there was no difference in the multiplication rate achieved at 1.07 μM NAA in presence of G-1, and the control treatment (autoclaved). When different concentration of PB were assayed together, at the best BA (6.99 μM) and NAA (1.07 μM) concentrations, the highest multiplication rate was achieved at the maximal PB concentration in presence of G-1 (Figure 1).

Immersion frequency in the proliferation and elongation phases

A 3-h immersion frequency significantly increased the multiplication rate relative to lower frequencies (Figure 2). Shoots grown in the TIB with a 1-h immersion frequency were taller and had a higher number of leaves, leaf area, and fresh and dry mass and total chlorophyll content than did shoots grown at longer immersion frequencies. All parameters above except the number of leaves per shoot were reduced for shoots grown at a 5-h immersion frequency (Table 1). Photosynthetic CO₂ assimilation of shoots in the 1 and 3-h immersion frequency treatments were not significantly different but were significantly higher than results for shoots grown in the TIB under the 5-h frequency (Table 1).

Table 1. Effect of immersion frequency on the growth parameters, net photosynthesis and chlorophyll content of pineapple shoots after 21 days in the culture previous acclimatization phase.

| Frequency | Shoot mass, (g) | | Shoot leaf area (m ²) | Total Chlorophyll (mg g ⁻¹ FW) | Net Photosynthesis (μmol CO ₂ m ⁻² s ⁻¹) | Shoot length (cm) | Leaves/shoot |
|-----------|-----------------|---------|-----------------------------------|---|--|-------------------|--------------|
| | fresh | dry | | | | | |
| 1h | 0.66 a | 0.037 a | 0.0118 a | 160 a | 8.07 a | 9.37 a | 6.66 ab |
| 3h | 0.49 b | 0.026 b | 0.0080 b | 145 b | 8.99 a | 6.83 b | 6.03 b |
| 5h | 0.18 c | 0.015 c | 0.0038 c | 105 c | 4.56 c | 4.52 c | 7.20 a |

Values represent the means of three replicated samples. Values indicated by different letters within columns are significantly different at the 5 % level by Tukey's Multiple Range Test.




Figure 1. Effect of plant growth regulator (PGR) concentration with Vitrofur G-1 sterilant on pineapple shoot multiplication rate after 30 days of proliferation in a temporary immersion bioreactor (TIB). The bar indicates the rate for autoclaved samples at the optimum PGR concentration. Data points ($n=3$ TIB samples) associated with the same letter were not significantly different as determined by the Tukey Test ($p < 0.05$).




Figure 2. The effect of immersion frequency in a temporary immersion bioreactor (TIB) on pineapple shoot multiplication rate. Bars having associated with the same letter were not significantly different ($p < 0.05$) as determined with the Tukey Test. Data are means of 3 TIB samples.

Discussion

Among the components of a culture medium, plant growth regulators are the best documented, although probably the least understood, elements controlling the morphogenetic response in micropropagation. The composition of plant growth regulators in the pineapple culture medium had as precedent the results obtained by Daquinta and Benegas (1997) in the propagation of 'Smooth Cayenne' using liquid medium. Later, Escalona *et al.* (1999) established that a supplement of $4.44 \mu\text{M}$ PB to this medium was optimal for pineapple proliferation in TIB.

Protocols for pineapple micropropagation generally use BA as the cytokinin, at concentrations between 2 and 5 mg L^{-1} . Using these concentrations, ten to 15 plants can be produced per month (Smith 2003). The role of BA to induce axillary proliferation in pineapple micropropagation is well documented (Smith 2003). During pineapple proliferation in TIB, when BA is reduced ($6.99 \mu\text{M}$) in the presence of PB ($4.44 \mu\text{M}$) the higher multiplication rate is achieved when NAA is omitted. However the use of $1.07 \mu\text{M}$ NAA permitted similar multiplication rate to control treatment and improved the length and shoot quality (Daquinta and Benegas, 1997).

When BA concentrations were tested in the presence of G-1, it was possible to achieve a higher multiplication rate at a lower BA concentration than was used when the medium was autoclaved. This result indicates that high temperatures during autoclaving affected the morphogenetic response of pineapple shoots in TIB. In this case, high temperatures could be denaturing proteins that can act as receptor sites at the membrane level.

The positive effect of paclobutrazol on pineapple micropropagation was previously described by Escalona *et al.* (1995) and was once again confirmed using TIB (Escalona *et al.*, 1999). Paclobutrazol controlled shoot growth and induced axillary bud proliferation. In the TIB, the use of paclobutrazol for pineapple micropropagation promoted formation of compact bud clusters with limited leaf development, avoiding unnecessary leaf growth during the shoot formation stage. It has been demonstrated that the use of inhibitors of gibberellin biosynthesis in tissue cultures containing a cytokinin increases the number of buds, especially with monocots. This phenomenon is not observed when media are supplemented with growth retardants only, indicating that they are not cytokinins per se. The interaction between gibberellins and growth retardants such as PB, is well documented. In *Spathiphyllum* plants, endogenous gibberellins seem to alleviate the shoot induction capacity of exogenous cytokinin. Imizadoles such as prochloraz inhibit the biosynthesis of gibberellins and as a consequence cytokinins can manifest their full shoot induction potential (Werbrouck *et al.*, 1996). The high multiplication rate we achieved when PB was used with reduced levels of BA and NAA in presence of G-1 could corroborate this statement.

In this study, we demonstrated that G-1 could be present in the medium continuously without seriously affecting pineapple shoot proliferation when BA and NAA levels present in the traditional medium established by Escalona (1999) are reduced. The main reason for this is probably that it combines ventilation of the plant tissues and intermittent contact between the entire surface of the tissue and the liquid medium (Etienne & Berthouly, 2002). The immersion frequency determined in this paper varies considerably between shooting and elongation phase. A high multiplication rate was achieved with an immersion frequency of 3h, while a one-hour immersion frequency gave shoots of a better quality, indicating that nutrient uptake is increased in this phase. The 1 h immersion

frequency using G-1 during the elongation phase increased not only morphological parameters of the shoots, but also the total chlorophyll contents and net photosynthesis. The photosynthetic capacity of shoots measured on day 21 of the elongation phase was higher than results obtained by Escalona *et al.*, (2003) at a 3 h immersion frequency without G-1. This factor constitutes an innovative step to take account during pineapple propagation in TIB.

The new procedure of pineapple propagation in the Temporary Immersion Bioreactor using G-1 introduces significant changes to the pineapple propagation scheme established by Escalona *et al.* (1999). First, it was possible to achieve a high multiplication rate at reduced concentrations of BA and NAA in the proliferation medium. Second, shoot elongation was most rapid at an immersion frequency of only 1h, which means the immersion frequency was increased to 24 times per day.

References

- Alvarado, Y., Herrera, L., Sarduy, Y., Acosta, M., Perez, B., and Carrazana, D. 1997. Actividad antimicrobiana del G-1 frente a bacterias en hongos contaminantes en la micropropagación de *Solanum tuberosum* L. tercer Seminario Internacional de Sanidad vegetal. Palacio de las Convenciones. Cuba.
- Daquinta, M. and Benegas, R. 1997. Brief review of tissue culture of pineapple. Pineapple News. (3): 7-9.
- Escalona, M., Lorenzo, J.C., González, B., Daquinta, M., Borroto, C., González, J.L. and Desjardins, 1999. Y. Pineapple micropropagation in temporary immersion systems. Plant Cell Rep. 18:743-748.
- Escalona, M., Lorenzo, J.C., Espinosa, P., González, B., Luna, I., Fundora, Z., Cintras, M. and Hernández, Z. 1995. Effect of paclobutrazol on *Ananas comosus* (L.) Merr. shooting. Paclobutrazol concentration. In: M. Estrada; E. Riego; J. Limonta; P. Téllez; J de la Fuente (Eds.) Advances in Modern Biotechnology. Vol III. II.4. ELFOS Scientae, La Habana, Cuba
- Escalona, M., Samson, G., Borroto C. G; Desjardins, Y. 2003. Physiology of effects of temporary immersion bioreactors on micropropagated pineapple plantlets. In Vitro Cell. Dev. Biol. Plant 39:651-656.
- Etienne, E. and Berthouly, M. 2002. Temporary Immersion Systems in plant micropropagation. Plant Cell Tissue Organ Cult. 69 (3): 215-231.
- Firoozabady, E. and Gutterson, N. 2003. Cost-effective in vitro propagation methods for pineapple. Plant Cell Reports 2-8.
- Fundora, Z., Cid, M., Pina, D., Escalona, M., Frómeta, Mosqueda, O. Efecto del G-1 en la propagación de la piña (*Ananas comosus* L (Merr) variedad cayena lisa BioVeg 2005. En Libro de Reportes Cortos Bioveg 2005: 102.
- George, M.W., Tripepi, R.R. 2001. Plant Preservative Mixture™ can affect shoot regeneration from leaf explants of chrysanthemum, European birch and rhododendron. HortScience 36:768-769.
- Murashige, T. and Skoog, F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. Physiol. Plant. 15: 473-497.
- Niedz, R. 1998. Using isothiazolone biocides to control microbial and fungal contaminants in plant tissue cultures. Hortechology 8:598-601.
- Porras, R.J. 1991 Recent advances and re-assessments in chlorophyll extraction and assay procedures for terrestrial, aquatic and marine organisms including recalcitrant algae. In: Scheer, H., ed. Chemistry of Chlorophyll. CRC Press Inc, Boca Raton, Ann Arbor, Boston, London; 320.
- Sestak, Z., Catsky, J. and Jarvis, P. G. 1971. Plant Photosynthetic production. In: Junk, W., ed. Manual of Methods. The Hague;. The Netherland N. V. Publisher;; 519.
- Smith, M.K., Ko, K.L., Hamill, S.D., Sanewski, G.M. and Graham. M.W. 2003. Biotechnology. In: Bartholomew D.P., Paul. R.E. and Rohrbach. K.G. (eds): The pineapple. Botany, production and uses. Pp 57-69.
- Werbrouck, S.P.O., Reding, P., Van Oncklen, H.A. and Debergh, P.C. 1996. Gibberellins play a role in the interaction between imidazole fungicides and cytokinins in Araceae. Journal of Plant Growth Regulation 14:105-107.◆

News From France

Growth Indicators for Queen Victoria Pineapple Versus Sum of Temperatures, Basis for a Heat-unit Model of Vegetative Growth

C.Dubois, P.Fournier, A.Soler, Cirad-Persyst alain.soler@cirad.fr

List of abbreviations : SC : Smooth Cayenne, WI : plant Weight Increase, WpW : Whole plant Weight, PIW : Weight of Planting material, DaP : number of Day after Planting.

Introduction

Queen Victoria is the main pineapple variety grown in Reunion Island which shows very clear-cut natural conditions (meaning of foregoing not clear). The pineapples are usually grown from sea level to an elevation of 900 m, with average annual temperatures varying from 19°C to 25°C, and annual rainfall from 1,500 mm to 5,000 mm. In these conditions, it is relatively easy to set up experiments to study the relationships between climatic parameters as temperatures and the growth of the plants.

Plant weight and D leaf weight are classical growth indicators for pineapple. The date of forced induction of flowering is usually determined in relation to the weight of the 'D' leaf (which is closely linked to plant weight, and therefore to fruit weight. Fleisch and Bartholomew (1987) and Malezieux et al. (1994 and 2003) built a heat-unit model able to predict the date of harvest. Heat-unit models have been developed to explained the impact of temperature on many aspects of plant growth and quality (Ganry, 1978; N'Da Adopo et al., 1996; Roman et al., 2000; Marzurczyk et al., 2003; Borreani et al, 2003; Tixier et al., 2007).

The aim of this study was to analyze the growth rate of Queen Victoria, and then to establish the basis to build a growth heat-unit model. First, changes in plant weight were monitored from planting to forcing, then changes in 'D' leaf weight were compared with the plant weights to determine the optimal value to be reached before forcing. Finally we looked for a relationship between growth of Queen Victoria and sum of temperatures.

Materials and methods

The trials were located at 21°19'21" E and 55°29'20" S, at an altitude of 150 m and were conducted on brown andic soils (pH=5.5). The pineapple was planted through plastic mulch and was provided with drip irrigation. For modelling purpose, two series of trials was set up in Reunion Island in 2007-2009 with Queen Victoria suckers. Suckers of comparable weight were planted in 2 different seasons, September (250 g) and February (225 g). Average temperature was lower, 21.8 °C, for the February planting, compared to 24.1 °C for September. Temperatures were recorded every 15 minutes with a data logger and the growth indicators (WI : plant Weight Increase, WpW : Whole plant Weight, D-leaf weight), were monitored on samples of 15 plants per observation.

Results - discussion

Increase in plant weight

The curve representing the increase in whole pineapple plant weight traditionally shows a sigmoid pattern (Py et al., 1984). However, in our case and during the observation periods (up to 230 days after planting), the curve showed a parabola pattern (Figure 1). The plant weight increase (WI) could be calculated as: $WI = WpW - PIW$ (where WpW = Whole plant weight, and PIW = Weight of Planting material). Then the plant weight increase (WI) could be expressed as a function of time (number of days after planting) : $WI = a \times DaP^2$ (where DaP = number of Days after Planting, and a is a calculated coefficient of the parabola curve).

'D' leaf weight

As previously observed for 'Smooth Cayenne' (Py et al., 1984), the 'D' leaf growth showed a sigmoid pattern with the following peculiarity (Figure 2): Weight increase of the 'D' leaf could be estimated by the equation (Verhulst, 1845): $y = y_{max}/(1+e^{-\hat{a}x})$ (where y_{max} represents the maximum weight reached by the 'D' leaf and \hat{a} and \hat{a} are calculated coefficients of the sigmoid curves). The maximum D leaf weight reached in our experimental conditions was 70 g for this variety. Exportable-size fruits (0.7kg) were obtained with a plant weight of about 1.2 kg at forcing with a 'D' leaf weight of 45 to 50 g. The plant weight for forcing (1.2 kg) was reached 210 to 230 days after planting under our experimental conditions.




Figure 1. Queen Victoria plants were planted in February and September and 15 plants were harvested, washed, dried and weighed each month. Weight increases over time were fitted and for February, $WI = 0.0233 \times DaP^2$ ($R^2 = 0.89$) and for September, $WI = 0.0272 \times DaP^2$ ($R^2 = 0.89$). Weight increases for the two dates were significantly different ($P < 0.05$).




Figure 2. Increase in D leaf weights of Queen Victoria pineapple plants. The data for February were fitted by the equation $D \text{ leaf weight} = 54.51 / (1 + \exp(2.27 - 0.02 \times DaP))$ - $R^2 = 0.61$; weights for September were fitted by the equation $D \text{ leaf weight} = 76.13 / (1 + \exp(3.48 - 0.0259 \times DaP))$ - $R^2 = 0.84$.

The growth (WI) of Queen Victoria was then related to the sum of temperatures and compared in each situation. For this preliminary stage, we added the daily averages of temperature above a threshold temperature for plant growth set at 8.34°C (Determined by successive approaches to maximize the coefficient of correlation).

During the 2 periods, it appeared that growth was linearly linked to temperature (Figure 3), and was also statistically identical, revealing the possibility of designing an original and relatively simple model in which pineapple growth is a function of temperature.




Figure 3. Growth in fresh weight of Queen Victoria as a function of sum of temperatures. Fifteen suckers were harvested monthly and were washed, dried and weighed. Temperature sums were accumulated by subtracting the threshold temperature of 8.34 °C from the daily mean temperature and summing. Equations for the lines were, for February, $WI = 0.6161 \times TTS - 742.7648 - R^2 = 0.65$, and for September, $WI = 0.5561 \times TTS - 622.6384 - R^2 = 0.70$ (differences not significant at $p > 0.05$).

Conclusion

At the current state of our observations, it appears that an increase in Queen Victoria plant weight is highly correlated with the sum of temperatures (linear relationship) in no-restrictive growth conditions, thus allowing us to establish the basis of a thermal time model of pineapple growth. Further data analysis for other varieties would indicate if this thermal time model may be a general model for all cultivated pineapples. Such a model is the basis for the development of more sophisticated models to design new sustainable cropping systems (Tixier et al., 2008).

References

- Borreani G., P.P. Roggero, L. Sulas, and M.E. Valente. 2003. Quantifying morphological stage to predict the nutritive value in Sulla (*Hedysarum coronarium* L.). *Agron.J.* 95:1608-1617.
- Fleisch, H., and D. Bartholomew. 1987. Development of a heat unit model of pineapple ('Smooth Cayenne') fruit growth from field data. *Fruits* 42:709-715.
- Ganry J. 1978. Recherche d'une méthode d'estimation de la date de récolte du bananier à partir de données climatiques dans les conditions des Antilles. *Fruits* 33:669-680.
- Malezieux, E., F.Cote, and D.Bartholomew. 2003. Crop environment, Plant Growth and Physiology. In *The Pineapple, Botany, Production and Uses*. CABI Publishing, Wallingford.
- Malezieux E., J. Zhang, E. Sinclair, and D. Bartholomew. 1994. Predicting pineapple harvest date in different environments, using a computer simulation model. *Agron. J.* 86:609-617.
- Mazurczyk W., B. Lutomirska, and A. Wierzbicka. 2003. Relation between air temperature and length of vegetation period of potato crops. *Agricultural and Forest Meteorology*, 118, p169-172.
- N'Da Adopo A., A. Lassoudière, and J. Tchango Tchango. 1996. Importance du stade de récolte pour la commercialisation de la banane plantain au Cameroun. *Fruits* 51:397-406.

- Py, C., J.J. Lacoëuilhe, and C. Teisson. 1984. The pineapple, cultivation and uses. Ed Maisonneuve et Larose, Paris. pp 568.
- Roman E.S., S.D. Murphy, and C.J. Swanton. 2000. Simulation of *Chenopodium album* seedling emergence. *Weed Science* 48:217-224.
- Tixier P., M. Dorel, and E. Malézieux. 2007. A model-based approach to maximize gross income by selection of banana planting date. *Biosystems engineering* 96:4/1-46.
- Tixier, P., E. Malézieux, M. Dorel, and J. Wery. 2008. SIMBA, a model for designing sustainable banana-based cropping systems. *Agricultural Systems* 97:139-150.
- Verhulst P.F. 1845. Recherches mathématiques sur la loi d'accroissement de la population. *Nouveaux Mémoires de l'Académie Royale des Sciences et Belles-Lettres de Bruxelles*. 18 : p. 1-42.

A Bait and Trap Method for Sampling Symphylid Populations in Pineapple

P.A. Marie-Alphonsine, P.Fournier, B. Dole, J.C. Govindin, A. Soler CIRAD-PERSYST. Corresponding author : A.Soler (alain.soler@cirad.fr).

Symphylids (*Hanseniella* sp) are polyphagous soil-borne parasites. On pineapple, they feed on roots tips (meristem tissues) producing either short roots with irregular swelling or roots with multiple branching resulting in the typical "witches brooms", (Kéhé, 1988; Py et al.1984; Petty et al., 2002). This behavior alters the root efficiency. The reddish plants look wilted with poor anchorage and growth. At field level, the crop appears as an irregular patchwork of diseased and normal plants, similar in appearance to nematode-induced symptoms (Py et al.1984; Kéhé, 1995; Soler, 1998; Perrier et al., 1993). Actually, diagnosing this disease in pineapple fields is rendered difficult due to possible confusion with other problems as water or nutrient deficiencies, nematodes, wilt (Kéhé, 1988; Py et al.1984).

As a consequence of the reduction in the use of pesticides due to environmental concerns, the control of symphylid populations in pineapple now requires particular and novel attention by researchers looking for an alternative to the chemical control of symphylid populations. Symphylids can be trapped and counted using appropriate bait or soil samples (Umble and Fischer, 2003a and Umble et al., 2006). We developed a reliable method to monitor symphylids in pineapple that combines bait and soil sample.

Material and methods

In a preliminary experiment we found that the trapping of symphylids was strongly improved by adding potato bait to the soil sample, compared to the soil sample alone or bait trapping alone. The bait and trap devices (Figure 1) are placed into the soil, carefully geolocalized and collected 3 days later for extracting and counting. Two methods for extracting and counting the symphylids were compared. In the first one (Figure 2A), soil aggregates were carefully broken apart before collecting the symphylids following the procedure of Umble (2006). In the second "floating" method (Figure 2B), the soil sample was carefully disaggregated in a 10 L bucket of water. Then floating symphylids were picked up with a small soft brush. The vertical distribution of the symphylid population was evaluated in pineapple and grass fallow rhizospheres (five replicates at four different depths from 4cm to 50cm).

Statistical analysis developed for nematode populations (Ferris, 1984, Perry et al., 2006) could, to some extent, be applied to symphylid populations, given that both are soil borne parasites developing in similar conditions and having many common traits (host-parasite relationship through the root system, pattern of development at field level, sensitivity to agro-climatic conditions). The spatial distribution of the populations in the field plots was evaluated using the "variance/mean" ratios (for details on calculations see Ferris, 1984, Perry et al., 2006) and spatial analyses based on Moran's and Geary's indices (for detailed calculations see Fortin, 1999 and Judas, 2002).

The method was tested in field plot of *Mucuna pruriens* cv *utilis* (a rotation crop) and in a grass fallow with different levels of infestation (plot areas 800m² with 60 samples randomly chosen). Then, the method was also tested on pineapple (cv MD-2) at different stages of development of the crop (plot of 50m² each with 5 samples randomly chosen and repeated several times over different seasons). Finally, the method can be used to evaluate rotation crop for pineapple (experiment in course).

Results and Discussion

Extraction methods

The floating method was more efficient in recovering juveniles (12.9 versus 7; $p < 0.001$) while recovery of total symphylids did not significantly differ between the two methods (Figure 3). Extraction durations were about six minutes per sample with both methods. The classical manual method allowed the recovery of symphylids in much better conditions for subsequent use (rearing for example) as the mortality was only 7% compared to 59% with the floating method. Nevertheless, as the adults/juveniles ratio can be used to determine if the population is growing or stagnant, the floating method appears to give a better picture of the evolution of the population.

Distribution of symphylids in soil profiles

Both tested sites, a grass fallow and a pineapple field, proved to be highly infested with an average of 47.2 and 44.6 symphylids respectively trapped in the grass fallow top soil and in the pineapple field. The number of symphylids did not significantly differ whatever the depth in both fields, implying that the infestation can spread not only at the surface but also deeper in the soil. Trapping symphylids only in the top soil (15cm) may be sufficient to monitor the population providing that climatic conditions at the time of sampling do not lead to a downward movement of the symphylids to find suitable conditions for their survival, for example during dry periods (Kéhé, 1988; Py et al.1984).

Examples of experiments and results allowed by the method

1) Maps of smoothed abundance data of symphylids observed at irregular locations (Figure 4) allowed a good visual evaluation of the distribution of the populations of symphylids in the field plots. Spatial analyses were made with the data obtained with different types of sampling (number of samples and number of traps per sample). The results showed that the symphylid populations are highly aggregated and that the range area for the development of symphylid populations appeared to be 4 to 6 meters wide, suggesting an optimized sampling with a regular spacing of about 4 meters between the traps ("variance / mean" ratios and Moran's and Geary's indices, Soler et al., submitted to Plant Dis.).

2) We used the bait and trap devices to monitor symphylid populations on a pineapple plot. Figure 5 shows that under pineapple, there were no symphylid up to the 4th month, then the populations increased up to forcing then declined after harvest. This probably reflects first, the efficiency of ethoprophos applied at planting and second, the reduced rhizogenic activity after harvest with no treatment since planting. The large standard deviations observed reflect the difficulty involved in obtaining a 'correct' sample in our small pineapple plots (50m²). But it also shows that the usual statistics (mean and standard deviation on raw data) are not very convenient for the description of aggregated populations of symphylids, as is also the case for nematode populations (Quénéhervé and Ferris, 1989).

3) The method is presently used to evaluate possible rotation crops as potential hosts of pineapple pests. Non-host crops could be used as rotation crops in new pineapple cropping systems (Figure 6). This experiment is presently in course.

Conclusion

Symphylids are one of the major pineapple pests in many production areas and their control has mainly been based on pesticide applications. With the reduction in the use of such pesticides due to environmental concerns, the control of this pineapple pest requires alternative to chemical applications. These alternative solutions will use ecologically-based integrated pest management (IPM). The method for trapping and counting symphylids described in this paper is an efficient tool to monitor symphylid populations in pineapple field plots and may help in validating new solutions for their control. Using this method we demonstrated symphylid populations are strongly aggregated, showing a range area for their development of about 4 to 6 meters and that they may be distributed homogeneously deep in the soil. Nevertheless, researchers need to take into account some basic facts concerning symphylid ecology, particularly their sensitivity to short-term climate variations such as drought or very wet conditions, which may temporarily affect the presence of symphylids in the upper part of a soil profile.

Literature cited

- Ferris, H. 1984. Nematode Damage Functions: The Problems of Experimental and Sampling Error. *Journal of Nematology* 16(1):1-9
- Fortin, M.J. (1999) Spatial statistics in landscape ecology. *Landscape ecological analysis: issues and applications* (eds J.M. Klopatek and R.H. Gardner), pp. 253-279. Springer, New York.
- Judas, M., K. Dornieden, U. Strothmann. 2002. Distribution patterns of carabid beetle species at the landscape-level. *Journal of Biogeography*, 29, 491-508.
- Kéhé, M. 1988. *Hanseniella ivorensis* - Juberthie-Jupeau et Kéhé (1978), Symphyle, myriapode (scutigerellidae) et le dépérissement racinaire de l'ananas en Côte d'Ivoire. PhD thesis. Université d'Abidjan. 180p.
- Kéhé, M., P. Gnonhoui, A. Adikoko. 1995. Time course of infestation by *Hanseniella ivorensis* (Symphylid) and *Pratylenchus brachyurus* (Nematode) on pineapple in Côte d'Ivoire. *Acta Horticulturae* 425: 465-474.
- Perrier X., Lacoeyille J.J., Malézieux E. 1993. An expert system for pineapple disorder diagnosis. *Acta Horticulturae* 334: 197-204.
- Perry, R.N., M. Moens. 2006. *Plant nematology*. Edts Perry RN And Moens, M. CABI Publishing, Wallingford, p 447.
- Petty, G.J., G.R. Stirling, D.P. Bartholomew. 2002. Tropical fruit pests and pollinators. Eds JE Pena, JL Sharp, M Wysoki. CABI Publishing, Wallingford, p 157-195.
- Py, C., J.J., Lacoeyille, C. Teisson. 1984. The pineapple, cultivation and uses. Ed Maisonneuves et Larose, Paris, pp568.
- Quénéhervé, P., Ferris H. 1989. Use of geometric vs arithmetic mean in the prominence value index and diagram frequency vs abundance.. *Journal of Nematology*, 2:1., 580.
- Soler, A. 1998 *Anan'AS : 150 fiches techniques pour maîtriser les techniques culturales de l'ananas*. Ed CIRAD-BEPAC, ISBN 2-912929-02-04, financial support by EU.
- Umble, J.R., Dufour R., Fisher G., Leap J., Van Horn, M., 2006; *Symphylans : soil pest management options*. NCAT ATTRA publication. pp16.

Umble, J. R., and J. R. Fisher. 2003a. Influence of below-ground feeding by garden symphylans (Ceph-alostigmata: Scutigereillidae) on plant health. *Environmental Entomology* 32: 1251-1261.

Umble, J. R., and J. R. Fisher. 2003b. Suitability of selected crops and soil for garden symphylan (Symphyla, Scutigereillidae: *Scutigereilla immaculata* Newport) population development. *Journal of Applied Soil Ecology* 24: 151-163.



Figure 1. Bait and trap device : how to use it? Black PVC "trapping" pots (250 ml volume, 9 cm high x 11 cm wide) with 76 holes 1.3 cm in diameter, 16 of which were at the bottom were buried 15 cm in the soil (entire pot was covered with soil). Prior to burial, the pots were gently filled to a depth of 3 cm with soil removed from hole in which the pot was to be buried. Bait consisting of three potato slices 2.5 cm in diameter and 2.5 cm thick was placed on the soil, the pot was filled to the top with additional soil and then buried.




Figure 2. "Manual" (A) and "floating" (B) methods of sampling symphylids. In the manual method, soil aggregates are carefully broken apart before collecting the symphylids. In the floating method, the soil sample was carefully disaggregated in a 10 L bucket of water and floating symphylids were picked up with a small soft brush.




Figure 3. Comparative efficiency of the "Manual" and "Floating" methods of symphyliid extraction. Fifteen symphyliids were inoculated in 250 ml of soil with organic matter. Samples were stored for 2 months at 29 °C with distilled water added daily to control humidity. Samples were counted after two months. There were ten replicates for each method.




Figure 4. Maps of smoothed symphyliid abundance data for plots planted to *Mucuna pruriens* cv *utilis*, (left) and grass fallow (right). Map scales in metres and the legends show numbers of symphyliids per sample.




Figure 5. Variations in symphylid populations under MD-2 pineapple. Plots included 300 plants on 6 ridges giving 50m² plots. Five traps with bait were placed with a minimum space of 4 m between 2 traps. Simple samplings were made 6 times at different periods of the year for each stage of development of the pineapple plants. Forcing at 12 months and Post harvest at 18 and 24 months.



Figure 6. Evaluation of rotation crop for pineapple. Left : the trap with the potato baits is placed inside the plot. The traps are collected 3 days later and the operation is repeated during several times during the growth of the crop. Right : partial view of the experimental site with *Crotalaria spectabilis* in front and *Crotalaria juncea* behind.



News from Ghana

Pineapple and Carbon Emissions

Rob Moss, PMB KD11, Kanda, Accra, Ghana. Tel/Fax: +233-21-232376; cell: +233 (0)243316080; website: www.waffcompany.com

Our organizations, West Africa Fair Fruit and the Dutch importer Agrofair, are undertaking a project to estimate the carbon footprint of pineapple in a commercial supply chain from Ghana to Europe using the recently released international standard PAS2050. We are doing this with consultants from the British science and agriculture consulting organization, ADAS. The end point will be in a verifiable carbon footprint (to be certified by the UK's Carbon Trust) and a plan for experimenting on how to reduce emissions.

Soils emit greenhouse gases as a consequence of fertilizer application and of soil processes. Soil emissions of nitrous oxide can be a significant contributor to the overall emissions from a product life cycle. Therefore to be able to make our footprint calculations complete we need to understand how soils used for pineapple production behave with respect to fluxes of nitrous oxide. Commercial pineapple grown in West Africa for the European market is planted through plastic mulch to conserve soil water. Farms growing pineapples tend to use very large amounts of nitrogen fertilizer, but some of this is applied to foliage above the plastic mulch. However, understanding of the source and sink strength of African soils for greenhouse gases is based very few relevant data. As part of our footprinting project for pineapple we are looking for any data on how soils behave in conditions comparable to those in the pineapple zone of Ghana and in a high fertilizer input system with plastic mulch covered ridges. There is a need to understand the fate of the applied nitrogen fertilizer (partitioning between plant, soil, losses to atmosphere and by soil water drainage), the moisture status of the soil, the degree to which there are anaerobic conditions under the plastic mulch, soil temperature under the plastic mulch, and how these conditions favour nitrous oxide emissions. The rainfall is about 1000 mm per year in the production zone we are working in. We would be very grateful if any researcher or organization that has any insights on this could contact either Rob Moss (robmoss@waffco.org) or Jeremy Wiltshire (Jeremy.wiltshire@adas.co.uk).

Ed. Note: Rob Moss contacted me by email to ask if I had information they were seeking. I did not and invited them to contact the "pineapple community" directly for assistance. Please help if you can. D.B.◆

News from Malaysia

Handling of Fresh-cut Pineapple for Fresh Consumption

Latifah, Mohd Nor, Horticulture Research Centre, Malaysian Agricultural Research and Development Institute, P.O.Box 12301, 50774 Kuala Lumpur, Malaysia; E-mail: lmn@mardi.gov.my.

ABSTRACT

Pineapple is one of the popular fruits served in fresh-cut form. At ambient temperature and without protective treatments, cut pineapple deteriorates rapidly, resulting in the development off-flavour and off-odour within a day. Fresh-cut pineapple sustains substantial tissue injury during processing; the disruption of tissue and cell integrity often increases respiration rate, ethylene synthesis, enzymatic browning and development of physiological disorders with associated increases in rates of other biochemical reactions responsible for changes in colour (including browning), flavor, texture and nutritional quality (sugar, acid and vitamin contents). The damaged plant tissues also provide a nourishing medium for microbial survival and growth. Chemical treatments such as sodium chloride, calcium chloride and ascorbic acid were used to improve the taste, flesh firmness and overcome the browning problem that occurred in 'Josapine' cut pineapple. The pineapple was cut into spears by machine (Fig 1A, 1B) and packed into rigid polypropylene containers (Fig 2C, 2D). Oxygen absorbent was inserted into the package for quality enhancement, which allows fresh-cut pineapple to be stored for 2 weeks at 2°C, 1 week at 10°C and 2 days at 25°C. Such storage periods provide sufficient time for fresh consumption of fresh-cut pineapple at the market shelf. A successful export trial was conducted in 2006 by MARDI by air shipping fresh-cut pineapple to the Netherlands. The quality of the fresh-cut pineapple remained good even after 6 days on the retail shelf. In 2009, shipment by refrigerated truck to Singapore was also successful. During both export trials, good collaboration was obtained from the Malaysian Pineapple Industrial Board (MPIB), Department of Agriculture (DOA), Federal Agricultural Marketing Authority (FAMA), local fruit exporters and fruit importers at the importing countries (Netherlands and Singapore).



Figure 1A, machine for cutting pineapple spears; 1B, fresh cut spears ready for packing; 1C and 1D, polypropylene containers packed with fresh-cut pineapple.



News From Taiwan

RAPD Marker Assisted Selection of EMS Induced Pineapple (*Ananas comosus* (L.) Merr.) Mutants

Yun-Hao Chen and Ching-Cheng Chen, Dept. Horticulture, National Chung Hsing University, 250, Kuo-kuang Rd., Taichung City 402, Taiwan, R.O.C.; E-mail: ccc@dragon.nchu.edu.tw

Introduction

Chemical mutagens have become more important for crop breeding and gene functions studies in recent years. Cross-breeding generally is the first choice in plant breeding programs but in some cases mutations induced on in vitro shoots, followed by clonal propagation and field selection procedures have higher probability of success (Predieri, 2001). Induced mutations, which change only one or a few specific traits of a cultivar but still meet the requirements of both the fruit industry and consumers, can offer a vast potential for fruit breeding (Predieri, 2001).

Molecular markers are detectable in all tissues regardless of the developmental stages and are not affected by the environment, pleiotropic and epistatic effects (Agarwal et al., 2008). The objective of the present study was to test the potential of using RAPD markers to analyze the putative mutants of pineapple that were produced by treatment with ethyl methane sulfonate (EMS). The effect of pH values of EMS solutions on germination rate of dormant axillary buds excised from pineapple crowns was also tested, which could be used as an alternative method for mutation induction in pineapple.

Material and methods

Fresh crowns of 'Tainung 17' pineapple were used in the present study. After the leaves were removed, crown stems were soaked in solutions containing 0.4% (v/v) EMS and 2% (v/v) dimethyl sulfoxide (DMSO), which was added as a carrier agent (Omar et al.,

1989), at different pH values for 8 hours. The pH was adjusted to 6.0, 7.0, or 8.0 with citrate-phosphate buffer (0.2M Na₂HPO₄ mixed with 0.1M citrate). The pH value of non-buffered 0.4% EMS solution was 3.75. Untreated crowns were used as the control. The stems of four crowns were the replicates for each treatment. After soaking, stems were washed three times with distilled water followed by sterilization with 1% sodium hypochlorite solution for 20 minutes. About 20 dormant axillary buds from each stem were aseptically excised and then incubated in half strength MS (Murashige and Skoog, 1962) medium supplemented with 30 g L⁻¹ sucrose, 8 g L⁻¹ agar, 2 ppm 6-benzyladenine (BA) and 0.2 ppm 1-naphthalene acetic acid (NAA). The pH value of the medium was adjusted to 5.8 before autoclaving. Germination rates of dormant axillary buds were recorded four weeks after incubation.

In another set of experiments, solutions containing 0, 0.02, 0.04, 0.06, 0.08, 0.1% (v/v) EMS were micro-filtered (0.22 µm; Minisart, Sartorius) and added into sterilized half strength MS medium. Tainung 17' *in vitro* plantlets (5mm-10mm in length) were cultured on the EMS mutation induction medium for about two months. The plants were then transplanted into pots and hardened in a net house. Three plants were randomly selected from each treatment and the genetic dissimilarity of those plants was analyzed by RAPD-PCR with ten Operon RAPD primers, OPA-03, OPA-17, OPG-03, OPG-08, OPG-16, OPG-18, OPJ-04, OPJ-11, OPJ-13 and OPJ-19. Based on the RAPD results, the proximity of the selected plants was analyzed and a dendrogram was generated by UPGMA (unweighted pair group method with arithmetic averages).

Results and discussion

In the EMS-DMSO study at different pH values, the bud germination rate in the pH 6.0 treatment was 36.5% after 4 weeks of incubation, which was significantly higher than that in the un-buffered treatment (Table 1). Apparently, adjusting the pH of EMS solution to near neutral reduced the phytotoxic effects of EMS (Table 1). Mhatre and Rao (2002) reported that the number of multiple shoots of pineapple formed decreased as the treatment duration and concentration of EMS solution increased. As a result, to be effective and efficient, concentration and pH of the mutagen solution, adding a carrier agent such as DMSO, and duration of treatment, all should be taken into consideration. In addition, treating dormant buds of pineapple crowns has a benefit over *in vitro* shoots because the shoots generated from the apex of the dormant buds can become new plants with a different genetic composition if mutations occur.

Molecular marker-assisted selection is a breeding method which provides the potential for increasing selection efficiency by allowing for earlier selection and reducing the plant population size required for field selection (Khawaleet al., 2007). RAPD is one type of molecular marker and the basis of this technique is using short random oligonucleotide sequences for differential PCR amplification of genomic DNA (Williams et al. 1991; Agarwal et al., 2008). The main advantages of RAPD are that it is simple, cheap and there is no requirement for DNA sequence information. However, its reproducibility is lower and it is also difficult to analyze approximate fragments (Tingey and Tufo, 1993; Schlötterer, 2004). In the present study, the average genetic similarities between EMS treated plants and the control decreased as the concentration of EMS increased (Table 2). The proximity analysis (Figure 1) also showed that higher concentrations of EMS induced plants that were more distinct from the control. It seems that the genetic dissimilarities of EMS treated plants gradually increased when the EMS concentration increased. This indicates that EMS concentration plays a critical role in the efficiency of mutation induction in pineapples and RAPD is able to detect the difference in EMS induced mutants.

Some induced morphological changes of EMS treated pineapple progenies were found in the present study such as changes in chlorophyll and anthocyanin contents in leaves, elongation and swelling of roots, leaf length and width, and spine patterns on leaf margin, etc.. These mutagen-derived morphologic traits might be useful for further gene function studies. In addition, those EMS treated plants produced in this study have been transplanted in the field for further evaluation and possible selection. Flowering and fruit characteristics will be examined.

References

- Agarwal, M., N. Shrivastava, and H. Padh. 2008. Advances in molecular marker techniques and their applications in plant sciences. *Plant Cell Rep.* 27: 617-631.
- Khawale, R. N., V. Yerramilli, and S. K. Singh. 2007. Molecular marker-assisted selection of *in vitro* chemical mutagen-induced grapevine mutants. *Curr. Sci.* 92: 1056-1060.
- Mhatre, M. and P. S. Rao. Influence of physical and chemical mutagens on pineapple shoot cultures. 2002. *Pineapple News* 9: 10-11.
- Murashige, T. and Skoog, F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.* 15: 473- 497.
- Omar, MS, F. J. Novak, and H. Brunner. 1989. *In vitro* action of ethylmethanesulphonate on banana shoot tips. *Sci. Hort.* 40: 203-295.
- Predieri, S. 2001. Mutation induction and tissue culture in improving fruits. *Plant Cell Tiss. Org. Cult.* 64: 185-210.
- Schlötterer, C. 2004. Opinion: The evolution of molecular markers - just a matter of fashion? *Nature Rev. Genet.* 5: 63-69.
- Tingey, S. V. and J. P. Tufo. 1993. Genetic analysis with random amplified polymorphic DNA markers. *Plant Physiol.* 101: 349-352.
- Williams, J. G. K., A. R. Kubelik, K. J. Livak, J. A. Rafalski, and S. V. Tingey. 1991. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res.* 18: 6531-6535.

Table 1. Regeneration of axillary buds of 'Tainung 17' pineapple crowns after an 8 h soak in 0.4% EMS-2% DMSO solutions having different pH values.[†]

| EMS-DMSO solution pH | Regeneration (%) |
|--------------------------|------------------|
| Check (no EMS treatment) | 49.3±3.5 |
| 3.75 | 9.5±10.3 |
| 6.0 | 36.5±20.9 |
| 7.0 | 21.2±10.8 |
| 8.0 | 28.3±25.0 |

[†]Crowns were treated with the EMS-DMSO solutions for 8 hrs before the dormant buds were excised and cultured in vitro for four weeks. Data represents the results of buds from stems of four crowns.

Table 2. Average genetic similarities between pineapple plantlets grown in media containing various concentrations of EMS and the control based on the RAPD analysis.[†]

| Concentration (%) | Genetic similarity |
|-------------------|--------------------|
| 0.02 | 0.948 |
| 0.04 | 0.928 |
| 0.06 | 0.923 |
| 0.08 | 0.896 |
| 0.10 | 0.896 |

[†] Three plantlets were incubated in medium containing each of the different concentrations of EMS for about two months.




Figure 1. Dendrogram of EMS treated 'Tainung No.17' pineapple plantlets generated by UPGMA cluster analysis based on similarity matrix of RAPD analysis. CK represents the control plantlet while 0.02-1~3, 0.04-1~3, 0.06-1~3, 0.08-1~3 and 0.1-1~3 represent plantlets from media containing 0.02% ~0.1% EMS.

News From the United States

Role of Proteases During Ripening of Pineapple Fruit

David A. Christopher, Dept. of Molecular Biosciences Bioengineering, University of Hawaii at Manoa, Honolulu, HI 96822. E-mail: dchr@hawaii.edu.

In a paper recently published in *Plant Physiology*, Neuteboom, et al. (<http://www.plantphysiol.org/cgi/content/abstract/151/2/515/>) describe a new process for the regulation of a protease and its inhibitor during pineapple fruit ripening. Fruit ripening and senescence involve hydrolysis and softening of tissues and a dramatic loss of chlorophyll. Various cellulases and proteases participate in the process. The most abundant cysteine protease in green and ripe pineapple fruit is bromelain, which accumulates without an inhibitory propeptide. How is bromelain regulated over time to avoid damaging cells until the stage of fruit development when it is needed for proteolysis? Cystatins are potent inhibitors of cysteine proteases, however the bromelains are enigmatically recalcitrant to inhibition by plant and animal cystatins. The researchers examined a novel secreted pineapple cystatin (AcCYS1), which has a uniquely long Alanine and Glutamate-rich N-terminus absent in other cystatins. They show the N-terminus is required for complete inhibition of stem and fruit bromelain. After the pineapple cystatin is synthesized, it inhibits the bromelain until ripening begins. Cleavage of the N-terminus from AcCYS1, a process which is specific to ripening fruit, renders the AcCYS1 inactive as an inhibitor against fruit bromelain, but not stem bromelain, thereby significantly enhancing bromelain activity in ripe fruit.

Environmental Friendly Approaches for Managing Nematodes and Weeds on Pineapple

Wang, K.-H.¹, B.S. Sipes¹, C.R.R. Hooks². ¹University of Hawaii, Department of Plant and Environmental Protection Sciences, Honolulu, HI; ² University of Maryland, Department of Entomology, College Park, MD, U.S.A.

Introduction

Reniform nematodes (*Rotylenchulus reniformis*) can reduce pineapple yields by up to 38% in the plant crop and by 60% in the ratoon crop. Currently, pre-plant fumigation with 1,3-dichloropropene (Telone[®]) or metam sodium (Vapam[®]) is practiced commonly in Hawaii to manage plant-parasitic nematodes. Because of environmental health concerns, increasing fuel costs, and the risk of losing soil fumigants, alternatives to soil fumigants are needed. Soil solarization (Fig. 1) and cover cropping are viable non-chemical alternatives for managing soil-borne plant-parasitic nematodes. Previously, it was determined that sunn hemp (*Crotalaria juncea*) (Fig. 2) possesses multiple mechanisms in suppressing reniform nematodes (Wang et al., 2001). These mechanisms include being a poor host of reniform nematodes, delaying the development of female reniform nematodes, producing an allelopathic compound that is toxic to the nematode, and enhancing nematode-trapping fungi that could prey on the nematode (Wang et al., 2001). However, suppressive effects of a sunn hemp cover crop against reniform nematodes are not consistent (Wang et al., 2002). Therefore, experiments are being conducted to determine if integrating sunn hemp cover cropping with soil solarization could improve reniform nematode suppression.

One challenge in managing reniform nematodes is their capability to survive in dry conditions through "anhydrobiosis." Anhydrobiosis is a survival strategy whereby the nematode coils and survives months to years without feeding (Tsai, 1978). Nematodes in their active stage are generally easier to kill by various nematode control strategies than when they are in their quiescent stage. Common pineapple plantation practice includes deep sub-soiling and fallowing the field for a year before the next crop planting. This may allow nematodes to enter an anhydrobiotic stage, making them more difficult to be killed. In addition, solarization only heats the top 10 to 20 cm of the soil (Chellemi, 1997; Wang et al., 2006). Thus, solarization might not be efficient in suppressing reniform nematodes that move deeper in the soil. Therefore, the objectives of the current research were to examine the potential of integrating sunn hemp cover cropping and solarization: 1) as alternatives to soil fumigation for nematodes and weed management; and 2) in improving soil health conditions as compared to fumigation.

Materials and Methods

A field trial was conducted at the Whitmore Experiment Station to compare the impacts of preplant treatments: 1) sunn hemp (SH) planted for 6 months, 2) solarized (Sol) for 2 months between March 27 and May 20, 2009, 3) SH planted for 4 months followed by 2 months of solarization (SH+Sol), and 4) fallow with weeds (control, C). Pineapple slips were planted into the prepared plots. At 3 months after pineapple planting (MAP), soil samples were taken from a nearby commercial pineapple field planted on the same date but fumigated with Telone[®] (1,3-d) followed by intensive herbicide application (Sipes, 2000). The purpose of this was to compare the above treatments with standard plantation practices. Soils were sampled prior to treatment installment (Pi1), after sunn hemp incorporation (Pi2), at the end of solarization (Pi3), and at 3 months after pineapple planting (MAP), which was done on July 9, 2009. Ten 20-cm deep soil cores were sampled per plot and composited. Nematodes were extracted from 250 cm³ of soil by elutriation (Byrd

et al., 1976) and counted. Number of pineapple planting holes with weeds present was recorded at 2 and 3 MAP. Soil health was compared among treatments and plantation practice at 3 MAP using nematode community analysis as described in Wang et al. (2006).

Results and Discussion

Solarization performed during spring-early summer in Hawaii generated significant amounts of heat, achieving a maximum temperature of 55 °C for both Sol and SH+Sol in the 0-10 cm soil layer. However, maximum temperature in the 11-20 cm soil layer only reached 37 °C in the Sol treatment and 41 °C in the SH+Sol treatment. Sol accumulated 174 hours above 42 °C, the lethal temperature for reniform nematodes, in the top soil layer, while the SH+Sol treatment accumulated 135 hours. Lethal temperatures were not reached in the deeper layer. Fourteen hours at 42 °C is required to kill reniform nematodes (Wang and McSorley, 2008). At 3 MAP, Sol did not significantly suppress reniform nematodes (Fig. 3). Numbers of reniform nematodes were reduced below the economic threshold level (1 reniform/g soil, i.e. ~250 reniform/250 cm³ soil) in the SH and SH+Sol treatments as compared to the control (Fig. 3). However, SH+Sol did not improve the reniform nematode suppressive effect as compared to SH. On the other hand, SH+Sol suppressed weeds more efficiently than SH or Sol alone (Table 1), and reduced weed pressure to one-third that of the control.

Free-living nematodes are good bioindicators of soil health because they are ubiquitous and have diverse feeding behaviors and life strategies (Bongers, 1990; Ferris et al., 2001; Neher, 2001; Yeates et al., 1990) and respond quickly to environmental soil disturbances (such as tillage, chemicals, pollutants, etc). Abundance or ratios of nematode trophic groups (Neher, 2001), diversity and richness and nematode faunal analysis (Ferris et al., 2001) are good indicators of soil health conditions. A comprehensive nematode community analysis was performed to evaluate soil health. Only nematode richness (number of genera), diversity, ratio of fungivores to bacterivores (F/B) and enrichment index (EI) are presented. EI is obtained by calculating the weight abundance of opportunistic bacterivores that responded quickly to nutrient enrichment and, therefore, is an indicator of soil rich in nutrients. Soil collected from the commercial pineapple field (T) had much lower richness than any of the treatments (Fig 4A). Although diversity was not different among treatments at 3 MAP, the commercial field and control treatment ranked lowest in diversity, another indication of less healthy soil conditions (Fig 4B). Higher F/B ratios usually indicate soil communities that are dominated by fungal decomposition rather than bacterial decomposition. Domination of fungal decomposition often is associated with environmental stress conditions (Ferris et al., 2001; Neher, 2001). At termination of solarization (Pi3), the F/B ratios indicated that Sol temporarily created a stressful condition, but this disturbance dissipated at 3 MAP (Fig. 4C). Integrating SH+Sol slightly reduced F/B at Pi3, indicating sunn hemp planting reduced the stress condition created by Sol (Fig. 4C). Although not significant, the highest F/B ratios were in the commercial field and in the control treatment.

Table 1. Pre-planting treatment effects on percent of pineapple planting holes with weeds present after planting on July 9, 2009.

| Treatment | 6 Aug 09 | 3 Sept 09 |
|------------------------|----------|-----------|
| Control (fallow) | 9.22 a | 14.50 a |
| Sunn hemp | 4.76 ab | 9.08 a |
| Sunn hemp+Solarization | 2.62 b | 4.39 b |
| Solarization | 5.54 ab | 8.74 ab |

Means are averages of 4 replications. Means in a column followed by same letters are not significantly different as determined by the Waller-Duncan (k-ratio) t-test. Data were transformed by the expression $\log(x + 1)$ prior to analysis.

Based on EI, SH significantly enriched soil nutrients at Pi2 (Fig 5). Although EI was temporarily reduced at termination of Sol (Pi3), this disturbance dissipated at 3 MAP. The commercial field had a lower EI than all treatments in the study area at 3 MAP, indicating a nutrient depleted condition. D-leaf weight was collected at 6 MAP, but no difference was observed among treatments.

In summary, planting sunn hemp without solarization significantly reduced the reniform nematode population below the economic threshold level at 3 MAP. Solarization only increased soil temperature at the top 10 cm of the soil and did not suppress reniform nematodes as compared to the control. Sunn hemp followed by a two-month solarization slightly increased soil temperature in the deeper soil layer as compared to Sol alone. Sol results could be better if Sol was performed during mid-summer. However, SH+Sol suppressed weed densities better than SH or Sol alone. Planting of sunn hemp enhanced nematodes involved in soil nutrient enrichment while the commercial field had the lowest enrichment index. On the basis of this index, the SH+Sol treatment is a more environmentally friendly option than standard pineapple plantation practices in Hawaii.

Literature Cited

- Byrd, D. W. Jr., K.R. Barker, H. Ferris, C. J. Nusbaum, W.E. Griffin, R. H. Small, and C.A. Stone. 1976. Two semi-automatic elutriators for extracting nematodes and certain fungi from soil. *Journal of Nematology* 8: 206-212.
- Chellemi, D. O, S. M. Olson, D. J. Mitchell, I. Secker, and R. McSorley. 1997. Adaptation of soil solarization to the integrated management of soilborne pests of tomato under humid conditions. *Phytopathology* 87, 250-258.
- Ferris H, T. Bongers, and R. G. M. de Goede. 2001. A framework for soil food web diagnostics: extension of the nematode faunal analysis concept. *Appl. Soil Ecol.* 18, 13-29.

- Neher, D. 2001. Role of nematodes in soil health and their use as indicators. *J. Nematol* 33, 161-168.
- Sipes, B.S. 2000. Crop profile of pineapples in Hawaii. <http://www.ipmcenters.org/CropProfiles/docs/hipineapples.html>.
- Tsai, B.-Y. 1978. Anhydrobiosis of the reniform nematode: survival and soiling. M.S. Thesis, University of Hawaii at Manoa, Honolulu.
- Wang, K.-H., R. McSorley, N. Kokalis-Burelle. 2006. Effects of cover cropping, solarization, and soil fumigation on nematode communities. *Plant and Soil* 286: 229-243.
- Wang, K.-H., B.S. Sipes, and D.P. Schmitt. 2001. Suppression of *Rotylenchulus reniformis* by *Crotalaria juncea*, *Brassica napus*, and *Tagetes erecta*. *Nematropica* 31: 235-249.
- Wang, K.-H., B.S. Sipes, and D.P. Schmitt. 2002. Management of *Rotylenchulus reniformis* in pineapple, *Ananas comosus*, by intercycle cover crops. *J. Nematol.* 34: 106-114.
- Wang, K.-H., B.S. Sipes, and D.P. Schmitt. 2003. Intercropping cover crops with pineapple for the management of *Rotylenchulus reniformis* and *Meloidogyne javanica*. *J. Nematol.* 35: 30-47.
- Wang, K.-H., and R. McSorley. 2008. Exposure time to lethal temperatures for *Meloidogyne incognita* suppression and its implication for soil solarization. *J. Nematology* 40: 7-12.
- Yeates, G. W., T. Bongers, R. G. M. DeGoede, D. W. Freckman, and S. S. Georgieva. 1993. Feeding habits in soil nematode families and genera - An outline for soil ecologists. *J. Nematol* 25, 315-312.



Figure 1. Soil solarization accomplished using transparent, 25- μ m-thick, uv-stabilized, low-density polyethylene mulch for two months.



Figure 2. Sunn hemp, *Crotalaria juncea*, is a tropical leguminous cover crop that has nematicidal properties and is commonly used as green manure.




Figure 3. Number of *Rotylenchulus reniformis* nematodes recovered from soil 3 months after planting pineapple at Whitmore, Hawaii. Treatments are fallow control (C), sunn hemp (SH), SH + solarization (SHSol) and solarization (Sol). Data are means of 4 replications. No differences were found after analysis by one-way ANOVA but significant differences between SH or no SH ($P < 0.05$) were found using contrast analysis. The data were transformed by the expression $\log(x + 1)$ prior to analysis.




Figure 4. Nematode richness (A), nematode diversity (B), and the fungivores:bacterivores ratio (C) at the initiation of the experiment (Pi1), termination of sunn hemp cover cropping (Pi2), termination of soil solarization (Sol) (Pi3) and 3 months after planting of pineapple (3). Symbols represent fallow control (C), sunn hemp (SH), two month solarization (Sol), SH followed by Sol (SH+Sol) and samples from a commercial field treated with 1,3-d and herbicide (T). Treatment means followed by the same letter are not significantly different, $P \leq 0.05$, $n=4$.




Figure 5. Enrichment index (EI) at the initiation of the experiment (Pi1), termination of sunn hemp cover cropping (Pi2), termination of soil solarization (Pi3) and 3 months after planting pineapple planting (3). Symbols represent fallow control (C), sunn hemp (SH), two-month solarization (Sol), SH followed by Sol (SH+Sol) and samples from a commercial field treated with 1,3-d and herbicide (T). Treatment means followed by the same letter are not significantly different, $P \leq 0.05$, $n=4$.

Commonalities of Pineapple, Agaves, and Cacti

Park S. Nobel, Distinguished Professor Emeritus, Department of Ecology and Evolutionary Biology, University of California, Los Angeles, CA 90095-1606

It all began with a taste test! Neimiah Grew reported in 1682 on the special acidic (bitter) taste of succulent plants in the morning. The taste test was expanded by Benjamin Heyne, who reported in 1815 that various succulents tasted much more acidic at dawn than at dusk. This nocturnal acidity increase was further studied in members of the Crassulaceae, becoming known as Crassulacean acid metabolism (CAM). As you all know, the most famous member of the family is pineapple, *Ananas comosus*. This bromeliad is cultivated on approximately 900,000 hectares worldwide for its fruit.

The biochemical details of CAM took a while to be elucidated. World War II stimulated advances in experimental methodology and instruments. Isotopes, especially ^{14}C , became available for tracing metabolic pathways, and chromatographic techniques were greatly improved. Based on work by Harry Beevers, Stanley Ransom, and Meirion Thomas in England and Hubert Vickery in the United States, much of the biochemistry of CAM was understood by the mid-1950s. Yet gas exchange studies indicating the daily pattern of CO_2 uptake and water vapor release by CAM plants were not interpreted with respect to the acidity studies until the 1960s.

The key feature of CAM shared by pineapple, agaves, and about 99% of cacti is the nocturnal uptake of CO_2 . At night, CO_2 from the atmosphere is joined onto the three-carbon compound phosphoenolpyruvate (PEP), a step catalyzed by PEP carboxylase in the cytosol of mesophyll cells. This leads to malate, oxaloacetate, and other acids that are responsible for the aforementioned nocturnal increase in acidity. These acids are stored during the night in the large central vacuoles of mesophyll cells. During the next daytime, the C_4 acids are released from the vacuoles, decarboxylated, and the CO_2 thus produced intracellularly is fixed by ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) in the chloroplasts of the mesophyll cells. This latter pathway is the same as that used by C_3 plants, which comprise about 92% of plant species including crops such as wheat, rice and most fruits and vegetables. About 1% of plant species (e.g., maize [corn], sugarcane, sorghum, tropical grasses) use the C_4 pathway, while 7% use the CAM pathway, our primary focus here. Of the three photosynthetic pathways, only CAM plants have nocturnal stomatal opening and the accompanying nocturnal uptake of CO_2 .

To understand the benefits of the CAM pathway, we shift our attention from CO_2 to water vapor. In particular, the much lower temperatures prevailing at night when the stomata of CAM plants open lead to much less water vapor diffusing out of them compared with C_3 plants and C_4 plants, whose stomata open during the daytime. A useful analogy is the rapid drying of a wet towel during the daytime versus the slow drying of a wet towel at night. This is because the water vapor concentration at saturation rises nearly

exponentially with temperature. And the inside of a pineapple or agave leaf (or a cactus stem) is essentially saturated with water vapor. The much lower force driving water vapor out of the leaves for the lower temperatures at night compared with the daytime leads to a higher water-use efficiency (WUE), the ratio of CO₂ fixed to water transpired. Indeed, the WUE of CAM plants averages about four-fold higher than for C₄ plants and eight-fold higher than for C₃ plants under the same environmental conditions.

Another interesting parameter is biomass productivity, which varies with plant species and photosynthetic pathway. Using measurements up to 1990, when such studies were fairly common, the five highest biomass-producing species averaged 41 tonnes of dry mass hectare⁻¹ year⁻¹ for C₃ plants, 52 for C₄ plants, and 42 tonnes hectare⁻¹ year⁻¹ for CAM plants. Pineapple was among the five highest-producing CAM species, with 35 tonnes hectare⁻¹ year⁻¹, while two agave species averaged 40 and two cactus species averaged 47 tonnes hectare⁻¹ year⁻¹. Thus, despite having the extra steps involved in nocturnal uptake and processing of CO₂ CAM plants can still have substantial biomass productivity.

The CAM species cultivated on the most area worldwide is the prickly pear cactus, *Opuntia ficus-indica*. More than 2.1 million hectares are cultivated in over 20 countries, mostly for forage and fodder. The area for the cultivation of its fruits, which are increasing called cactus pears to avoid drawing attention to the nasty spines, is about 140,000 hectares. The country currently with the largest cultivation of *O. ficus-indica* is Brazil, with 600,000 hectares, with nearly as much cultivated in Tunisia, in both cases mainly for forage and fodder. The most extensively cultivated agave is *Agave tequilana*, with 80,000 hectares used for tequila production in Mexico, while about half as much area is devoted to other agave species for other beverages.

How will net CO₂ uptake and biomass productivity of pineapple, agaves, and cacti be affected by global climate change? With regard to increasing atmospheric CO₂ levels, the news is good, as productivity increases by about 1% for each 10 ppm increase in atmospheric CO₂ level (currently, the CO₂ level is increasing at just over 2 ppm per year). None of these three taxa love freezing temperatures, so the generally increasing global temperatures are also good news. In addition, CAM plants cope well with changes in rainfall patterns. All of these influences on net CO₂ uptake and productivity can be quantified using an Environmental Productivity Index (EPI).

EPI indicates the fraction of maximal net CO₂ uptake over 24-hour periods under any set of environmental conditions. It incorporates individual multiplicative indices for the effects of light, temperature, water status, nutrients, and atmospheric CO₂ level. EPI has been used to interpret and predict the productivity of agaves and cacti, and it is ripe for application to pineapple. More detail on EPI and the other topics presented above are described in P.S. Nobel, "DESERT WISDOM/AGAVES and CACTI: CO₂, Water, Climate Change" published by the author in 2010 and available through iuniverse.com, amazon.com, and barnesandnoble.com for \$16.95 (\$6.00 for the ebook available through iuniverse.com).

Ed. Note: Distinguished Professor Park Nobel is one of a few people who has spent much of a career investigating the productivity of plants with CAM. I had not thought of cacti as being important agricultural plants until I began reading his interesting book and found it to be most interesting. I think readers of **Pineapple News** would be interested to know of his work. I believe there are principles in the book that we "pineapple people" can learn from. ♦

Services

The listings under Commercial Services and Directory of Professionals is maintained as a convenience to readers and should in no way be construed as an endorsement of those providing commercial or professional services. Those offering specialized services to pineapple growers or researchers are invited to contact the editor for possible inclusion in the listings below.

Commercial Services

Maintain CF 125 continues to be available for use in pineapple plant propagation. A renewal letter for registration of the product was received in 2003. For further information, contact Bhushan Mandava, Repar Corporation, P.O. Box 4321, Silver Spring, MD 20914 Tel: 202-223-1424 Fax: 202-223-0141; E-Mail: mandava@compuserve.com

Centro de Bioplantas. Dr. Justo L. Gonzalez Olmedo, Director of Foreign Affairs Office, Centro De Bioplantas. Universidad De Ciego De Avila, Carretera a Moron Km 9. Cp69450. Cuba. Centro De Bioplantas offers certificates of authenticity for pineapple material propagated in their tissue culture facility. Web site: <http://www.Bioplantas.cu>

LAMERSA, Dole's meristem laboratory in Honduras. Contact John T. Mirenda PhD, Dole Fresh Fruit International Ltd., San Jose, Costa Rica. Phone: 506 287 2175. Fax: 506 287 2675. E-mail: Jmirenda@la.dole.com. The laboratory can produce meristematically-derived plants of pineapple as well as banana and other crops.

Thai Orchids Lab, Dr. Paiboolya Gavinlertvatana. Horticulture/ agriculture/ forestry tissue culture laboratory with exports to Australia, U.S.A., Africa, and Asia. MD2 pineapple available (open to acquiring additional varieties) or confidential exclusive contract propagation. Phone: +1 510 931 7865 Fax: +66 2510 9452 Website: <http://www.tolusa.com/> E-mail: info@tolusa.com.

Vitropic, Zone d'Activités Economiques des Avants, 34270 Saint Mathieu de Trévières France; Tel: + 33 (0)4 67 55 34 58; Fax: + 33 (0)4 67 55 23 05. E-mail : vitropic@vitropic.fr. Web site: www.vitropic.fr. Vitropic proposes the best individuals from the CIRAD FHLOR selected clones including: Cayenne Group, Queen Group, Perolera Group, MD2, Ornamentals pineapples. The range is continuously extending, do not hesitate to ask for more information.

Professional Services

Mr. Wilbert Campos Alvarado. M.Sc. Tropical Soils & Crop Mgmt. E-mail. wcamposa@gmail.com. Phone: (506) 8815-7271. Apdo. Postal 536-7210, Guapiles, Costa Rica. Experience in all stages of production (soil preparation, plant nutrition, diseases & pest control, PGR use, etc) of pineapple for the fresh fruit production market as well as experience in packing plant management and in postharvest treatment. Also worked in pineapple R&D for several years under different climate conditions (Costa Rica, Guatemala, Ecuador).

Ing. Alejandro Chavarría. APDO 4437-56 Pital, San Carlos. Alajuela, Costa Rica. Tel: (506) 88-20-79-55 / (506) 24-73-40-00, alechava@hotmail.com. I have worked like an International Pineapple Consulting in México, Costa Rica and Brazil. Experienced in project feasibility, plantation design, agricultural machinery, all aspects of farm crop management, post harvest management and establishment of good agricultural practices.

Dr. Mark Paul Culik. INCAPER, Rua Alfonso Sarlo 160, CEP 29052-010, Vitoria, ES, Brazil; Tel: 27-3137-9874; markculik3@yahoo.com. Experience: PhD in Entomology with more than 25 years of agricultural pest management experience in crops ranging from apples to papaya and pineapple, identification of pests and beneficial arthropods ranging from Collembola to fruit flies, and current work on scale insects with emphasis on pineapple mealybugs. Areas of specialization: Entomology, Insect and Pest Identification, Integrated Pest Management.

Dr. Francisco Gomez (E-mail: fgomez1@cablecolor.hn) and **Jose R. Vasquez**, MBA (E-mail: jrva46@excite.com). Golden Pacific Ag Services, PO.Box 15088, Lomas Miraflores, 4a. Calle, 1a Avenida # 4326, Tegucigalpa, Honduras. Phone: 504 230 1120; 504 969 5568. Experience: Pineapple and melon production, from seed propagation-planting-field maintenance-forcing-harvesting-post-harvest management and commercialization.

Mr. L. Douglas MacClure. 360 Hoopalua Dr., Pukalani, Hawaii, U.S.A. E-mail: norfolkldm@aol.com. Experience: More than 39 years with Maui Pineapple Company heading plantation and diversified agriculture operations and started the Royal Coast Tropical Fruit Company in Costa Rica. Collected and summarized production information in Asia and Central America. Also consulted on pineapple for companies and growers in El Salvador, Australia, Thailand and Indonesia.

Mr. Graham J. Petty 13 Somerset Place, Lambert Road, Port Alfred, 6170, Republic of South Africa. Phone: +27 (0) 46 624 4868; Tel/Fax: +27 (0) 46 625 0946; E-mail: grahamp@imaginet.co.za. Experience: M.Sc. (Agric) Pretoria : Pr. Sci. Nat. . Researcher and advisor to the South African Canning Pineapple Industry on matters of Pest Management in pineapple culture, for 34 years. Economic entomology and management of biological control agents have received particular attention.

Mr. Col Scott. E-mail: scottch45@bigpond.com. Mobile: +61 488092442; Phone: +61 7 34252417; Fax: +61 7 34252417. Over 37 years experience in all aspects of pineapple agronomy and research in Australia (32 years with Golden Circle Ltd) and South Africa (5 years with Summerpride Foods Ltd). Experience includes working with growers, researchers and fertilizer and agricultural chemical suppliers. Other production areas visited include Hawaii, Central America, Thailand, Indonesia and Malaysia.

Dr. José Aires Ventura. Incaper, Rua Afonso Sarlo 160 (bento Ferreira), 29052-010, Vitoria-ES, Brazil. E-mail: ventura@incaper.es.gov.br; Tel.: 55-27-31379874. www.incaper.es.gov.br. Area of Specialization: Plant Pathology (research in pineapple diseases management; Fusarium diagnosis, diseases resistance).

Mr. Dean Wheeler. AgResults Inc., 609 Buchanan Street, Davis, California, U.S.A. 95616. Phone/fax: 530-758-4620 Residence: 530-758-3354. Email: agresults@aol.com. Web page at <http://agresults.com/>.

Book Reviews and Web Sites

Book Reviews

No reviews were provided for this issue.

Web Sites of Possible Interest

Nothing new at this time.◆

New References on Pineapple

The list below includes papers related to various aspects of pineapple culture, physiology, processing, preservation or byproducts that were published or located since the last issue of the newsletter was printed. Some papers may seem relatively unrelated to pineapple but since judgement must be exercised when including or excluding references, the decision was made to err on the side of inclusion so as to serve as many readers as possible. Often, abstracts of the papers listed below can be found on-line and of course all abstracts of paper published in *Acta Horticulturae* are available from info@ishs.org.

- Abdullah, H., Rohaya, M.A., and Hasmah, E.A.E., 2008. Increasing pineapple fruit resistance to chilling injury during storage by temperature preconditioning. *Acta Horticulturae* 768:217-224.
- Abilio, G.M.F., Holschuh, H.J., Bora, P.S., and Oliveira, E.F.d., 2009. Extraction, bromelain activity and analysis of some chemical parameters in pineapple varieties from Paraiba. *Revista Brasileira de Fruticultura* 31:1117-1121.
- Agergaard, J., Fold, N., and Gough, K.V., 2009. *Rural-urban dynamics: livelihoods, mobility and markets in African and Asian frontiers*. Routledge, Abingdon.
- Ahmad, B., Rathar, G.M., Varshney, A., and Khan, R.H., 2009. pH-dependent urea-induced unfolding of stem bromelain: unusual stability against urea at neutral pH. *Biochemistry (Moscow)* 74:1337-1343.
- Ajibola, V.O., Babatunde, O.A., and Suleiman, S., 2009. The effect of storage method on the vitamin C content in some tropical fruit juices. *Trends in Applied Sciences Research* 4:79-84.
- Akali, S., Jha, K.K., and Amod, S., 2009. Cultivation of pineapple for upliftment of rural economy in Nagaland, India. *Acta Horticulturae* 822:99-106.
- Alavi, G., Sanda, M., Loo, B., Green, R.E., and Ray, C., 2008. Movement of bromacil in a Hawaii soil under pineapple cultivation - a field study. *Chemosphere* 72:45-52.
- Albornoz, A., Ortega, L., Segovia, E., Bracho, Y., and Cubillan, G., 2009. Purchase attributes of fresh fruits at domestic level. *Bioagro* 21:57-62.
- Almeida, C.O.d., Matos, A.P.d., Cardoso, C.E.L., Sanches, N.F., Teixeira, F.A., and Elias Junior, J., 2009. Evaluation of economic and environmental impacts of integrated pineapple production in the State of Tocantins-Brazil. *Acta Horticulturae* 822:161-168.
- Almeida, C.O.d., Matos, A.P.d., Cardoso, C.E.L., Sanches, N.F., Teixeira, F.A., and Elias Junior, J., 2009. Evaluation of social impact of integrated pineapple production in the State of Tocantins-Brazil. *Acta Horticulturae* 822:169-176.
- Almeida, C.O.d. and Reinhardt, D.H.R.C., 2009. Pineapple agribusiness in Brazil. *Acta Horticulturae* 822:301-312.
- Almeida, M.M.B., Sousa, P.H.M.d., Fonseca, M.L., Magalhães, C.E.C., Lopes, M.d.F.G., and Lemos, T.L.G.d., 2009. Evaluation of macro and micro-mineral content in tropical fruits cultivated in the northeast of Brazil. *Ciencia e Tecnologia de Alimentos* 29:581-586.
- Alothman, M., Rajeev, B., and Karim, A.A., 2009. Antioxidant capacity and phenolic content of selected tropical fruits from Malaysia, extracted with different solvents. *Food Chemistry* 115:785-788.
- Anonymous, 2008. *Pantoea citrea*. [Distribution map]. Map 1038.
- Antony, E., Taybi, T., Courbot, M., Mugford, S.T., Smith, J.A.C., and Borland, A.M., 2008. Cloning, localization and expression analysis of vacuolar sugar transporters in the CAM plant *Ananas comosus* (pineapple). *Journal of Experimental Botany* 59:1895-1908.
- Araujo, K.G.L., Sabaa-Srur, A.U.O., Rodrigues, F.S., Manhães, L.R.T., and Canto, M.W.d., 2009. Utilization of pineapple (*Ananas comosus* L.) cv. Perola and Smooth cayenne for wine making - study of chemical composition and acceptance. *Ciencia e Tecnologia de Alimentos* 29:56-61.
- Arun, D., Deb, P., Roy, A., and Suresh, C.P., 2009. Effect of foliar feeding of commercial formulations of growth regulators and micronutrients on some physico-chemical characters of pineapple (cv Kew). *Environment and Ecology* 27:114-116.
- Asrey, R., Patel, V.B., Singh, S.K., and Sagar, V.R., 2008. Factors affecting fruit maturity and maturity standards - a review. *Journal of Food Science and Technology (Mysore)* 45:381-390.
- Aziz, M.G., Hierro, A.M.d., Mazumder, M.A.R., Uddin, M.B., and Kulbe, K.D., 2009. Effect of enzymatic hydrolysis of pineapple fruit pulp on yield and analytical parameters of derived juice. *International Journal of Sustainable Agricultural Technology* 5:29-35.
- Balasundram, S.K., Husni, M.H.A., and Ahmed, O.H., 2008. Spatial variability of pineapple yields on a tropical peat. *Crop Management*:0418-0401-RS.
- Baldotto, L.E.B., Baldotto, M.A., Giro, V.B., Canellas, L.P., Olivares, F.L., and Bressan-Smith, R., 2009. Performance of 'Victoria' pineapple in response to humic acid application during acclimatization. *Revista Brasileira de Ciencia do Solo* 33:979-990.
- Bi, J., Fang, L., Ding, Y., Wang, P., and Bai, S., 2009. Optimization of explosion puffing drying for pineapple at variable temperature and pressure difference. *Transactions of the Chinese Society of Agricultural Engineering* 25:334-339.
- Borsato, D., Moreira, I., Nobrega, M.M., Moreira, M.B., Silva, R.S.d.S.F.d., and Bona, E., 2009. Modeling and simulation of the osmotic dehydration process of pineapple's pieces using the finite element method. *Quimica Nova* 32:2109-2113.
- Braga, A.M.P., Pedrosa, M.P., Augusto, F., and Silva, M.A., 2009. Volatiles identification in pineapple submitted to drying in an ethanolic atmosphere. *Drying Technology* 27:248-257.
- Brito Neto, J.F.d., Pereira, W.E., Sá;Sobrinho, R.G.d., Barbosa, J.A., and Santos, D.P.d., 2008. Productive aspects of the family and commercial pineapple culture in the state of Paraíba. *Caatinga* 21:43-50.
- Cabral, A.C.S., Said, S., and Oliveira, W.P., 2009. Retention of the enzymatic activity and product properties during spray drying of pineapple stem extract in presence of maltodextrin. *International Journal of Food Properties* 12:536-548.
- Cabral, J.R.S., Ledo, C.A.d.S., Caldas, R.C., and Junghans, D.T., 2009. Characters variation in pineapple hybrids obtained by different crosses. *Revista Brasileira de Fruticultura* 31:1129-1134.
- Cabral, J.R.S. and Matos, A.P.d., 2009. Imperial, a new pineapple cultivar resistant to fusariosis. *Acta Horticulturae* 822:47-50.

- Cabral, J.R.S., Matos, A.P.d., Junghans, D.T., and Souza, F.V.D., 2009. Pineapple genetic improvement in Brazil. *Acta Horticulturae* 822:39-46.
- Cambui, C.A., Gaspar, M., and Mercier, H., 2009. Detection of urease in the cell wall and membranes from leaf tissues of bromeliad species. *Physiologia Plantarum* 136:86-93.
- Carvalho, A.C.P.d., Pinheiro, M.V.M., Dias, G.d.M.G., and Morais, J.P.S., 2009. In vitro multiplication of ornamental pineapple by shoot etiolation and regeneration. *Horticultura Brasileira* 27:103-108.
- Chakraborty, A.J. and Bera, B., 2008. A study on structure of costs and returns of pineapple cultivation in Darjeeling district of West Bengal. *Journal of Crop and Weed* 4:24-30.
- Chan, Y.K., 2008. Breeding of seed and vegetatively propagated tropical fruits using papaya and pineapple as examples. *Acta Horticulturae* 787:69-75.
- Chang, H., Li, Y., and Yang, X., 2008. Investigation of the drying characteristics of pineapple powder. *Modern Food Science and Technology* 24:5-7.
- Chauhan, O.P., Alok, S., Asha, S., Raju, P.S., and Bawa, A.S., 2009. Modeling of pre-treatment protocols for frozen pineapple slices. *LWT - Food Science and Technology* 42:1283-1288.
- Chen, N.J., Paull, R.E., Chen, C.C., and Saradhulhat, P., 2009. Pineapple production for quality and postharvest handling. *Acta Horticulturae* 822:253-260.
- Coelho, R.I., Carvalho, A.J.C.d., Thiebaut, J.T.L., and Lopes, J.C., 2009. Bud sprouting in stem sections of 'Smooth Cayenne' pineapple plant treated with growth regulators. *Revista Brasileira de Fruticultura* 31:203-209.
- Corcho Fuentes, J.S.P.d., Pegna, F.G., Iglesias Coronel, C., Garcia Reina, F., and Spugnoli, P., 2009. Power demand of a flail mower during pineapple field demolition. *Ciencia e Investigacion Agraria* 36:59-68.
- Cordeiro, I.M.C.C., Santana, A.C.d., Lameira, O.A., and Silva, I.M., 2009. Economical analysis of cultivation systems with *Schizolobium parahyba* var. amazonicum (Huber ex Ducke) Barneby (Parica) and *Ananas comosus* var. *erectifolius* (L. B. Smith) Coppus & Leal (Curaua) crop at Aurora do Pará, Brazil. *Revista de la Facultad de Agronomía, Universidad del Zulia* 26:243-265.
- Correia, D., Rocha, M.V.P., and Alvez, G.C., 2009. Growth of micropropagated *Ananas comosus* var. *erectifolius* plantlets in different substrates under greenhouse conditions. *Acta Horticulturae* 822:85-89.
- Costa, H.B., Delboni, S.G., Fortunato, F.S., and Ventura, J.A., 2009. Proteolytic activity in stems of 'Vitoria', 'Smooth Cayenne' and 'Perola' pineapple plants. *Acta Horticulturae* 822:239-243.
- Cruvinel, R.S.R., Prado, M.E.T., Resende, J.V.d., and Cirillo, M.Ã., 2009. Preservation of frozen pineapple pieces through the vacuum impregnation with pectin and calcium propionate. *Revista Ciencia Agronomica* 40:539-548.
- Culik, M.P. and Ventura, J.A., 2009. New species of *Rhinoleucophenga*, a potential predator of pineapple mealybugs. *Pesquisa Agropecuária Brasileira* 44:417-420.
- Culik, M.P., Ventura, J.A., and Martins, D.d.S., 2009. Scale insects (Hemiptera: Coccidae) of pineapple in the State of Espírito Santo, Brazil. *Acta Horticulturae* 822:215-218.
- Dacera, D.d.M., Babel, S., and Parkpian, P., 2009. Potential for land application of contaminated sewage sludge treated with fermented liquid from pineapple wastes. *Journal of Hazardous Materials* 167:866-872.
- Damasceno e Silva, K.J., Souza, V.A.B.d., and Gomes, R.L.F., 2008. Effect of plantlet heights on post-in vitro-culture adaptation of ornamental pineapple. *Revista Ceres* 55:551-555.
- Dantas Junior, O.R., Silva, S.d.M., Alves, R.E., and Silva, E.d.O., 2009. Susceptibility to chilling injury for 'Perola' pineapple treated with 1-methylcyclopropene. *Revista Brasileira de Fruticultura* 31:135-144.
- Dhar, M., Rahman, S.M., and Sayem, S.M., 2008. Maturity and post harvest study of pineapple with quality and shelf life under red soil. *International Journal of Sustainable Crop Production* 3:69-75.
- Dionello, R.G., Berbert, P.A., Molina, M.A.B., Pereira, R.C., Viana, A.P., and Carlesso, V.O., 2009. Osmotic dehydration of sliced pineapple of two cultivars in inverted sugar syrup. *Revista Brasileira de Engenharia Agrícola e Ambiental* 13:596-605.
- Dionello, R.G., Berbert, P.A., Molina, M.A.B.d., Pereira, R.d.C., Viana, A.P., and Carlesso, V.d.O., 2009. Assessment of convective drying models for fresh and osmo-dehydrated pineapple rings. *Ciencia e Tecnologia de Alimentos* 29:232-240.
- Egidio, V.d., Sinelli, N., Limbo, S., Torri, L., Franzetti, L., and Casiraghi, E., 2009. Evaluation of shelf-life of fresh-cut pineapple using FT-NIR and FT-IR spectroscopy. *Postharvest Biology and Technology* 54:87-92.
- Elias Junior, J., Gomes, D.C., Matos, A.P.d., and Almeida, C.O.d., 2009. Micro and macroeconomic analyses of the pineapple industry in the state of Tocantins. *Acta Horticulturae* 322:317-322.
- Elzebroek, T. and Wind, K., 2008. Edible fruits and nuts, p. 25-131. CABI, Wallingford.
- Faria, D.C.d., Carvalho, A.J.C.d., Coelho, R.I., and Oliveira JÃºnior, J.C.d., 2009. Bud moth (*Strymon megarus* (Godart, 1824)) causes severe losses in young plants of pineapple cultivars 'Perola' and smooth cayenne in Rio de Janeiro, Brazil. *Acta Horticulturae* 822:219-224.
- Fawole, O.P., 2008. Pineapple farmers' information sources and usage in Nigeria. *Bulgarian Journal of Agricultural Science* 14:381-389.
- Ferreira, A.C.H., Neiva, J.N.M., Rodriguez, N.M., Campos, W.E., and Borges, I., 2009. Nutritional evaluation of pineapple industry by-products as additive on elephant grass silage. *Revista Brasileira de Zootecnia* 38:223-229.
- Ferreira, A.C.H., Neiva, J.N.M., Rodriguez, N.M., Santana, G.Z.M., Borges, I., and LÃ´bo, R.N.B., 2009. Performance of sheep fed with silages of elephant grass mixed with by-products from tropical fruit juices industry. *Revista Ciencia Agronomica* 40:315-322.
- Ferreira, E.A., Costa, F.H.S., and Pasqual, M., 2009. Acclimatization of micropropagated plantlets of 'Imperial' pineapple. *Acta Horticulturae* 822:91-95.
- Ferreira, E.H.d.R., Rosenthal, A., Calado, V., Saraiva, J., and Mendo, S., 2009. *Byssoschlamys nivea* inactivation in pineapple juice and nectar using high pressure cycles. *Journal of Food Engineering* 95:664-669.

- Fitzhugh, D.J., Shan, S.Q., Dewhirst, M.W., and Hale, L.P., 2008. Bromelain treatment decreases neutrophil migration to sites of inflammation. *Clinical Immunology* 128:66-74.
- Fold, N. and Gough, K.V., 2008. From smallholders to transnationals: the impact of changing consumer preferences in the EU on Ghana's pineapple sector. *Geoforum* 39:1687-1697.
- Fraguas, C.B., Dornelles, C.M.d.V., and Lima, G.P.P., 2009. In vitro bud induction and multiplication of cv. 'IAC Gomo-de-mel' pineapple fruit with benzyl amino purine and naphthalene acetic acid. *Ciencia Rural* 39:1682-1687.
- Freschi, L., Nievola, C.C., Rodrigues, M.A., Domingues, D.S., Sluys, M.A.v., and Mercier, H., 2009. Thermoperiod affects the diurnal cycle of nitrate reductase expression and activity in pineapple plants by modulating the endogenous levels of cytokinins. *Physiologia Plantarum* 137:201-212.
- Gangopadhyay, G., Mukherjee, K.K., Gangopadhyay, S.B., and Roy, S.K., 2009. Agrobacterium-mediated genetic transformation of pineapple var. Queen using a novel encapsulation-based antibiotic selection technique. *Plant cell, tissue, and organ culture* 97:295-302.
- Garcia, R., Alves, E.S.S., Santos, M.P., Aquije, G.M.F.V., Fernandes, A.A.R., Santos, R.B.d., Ventura, J.A., and Fernandes, P.M.B., 2008. Antimicrobial activity and potential use of monoterpenes as tropical fruits preservatives. *Brazilian Journal of Microbiology* 39:163-168.
- Gaurab, G., Roy, S.K., Gangopadhyay, S.B., and Mukherjee, K.K., 2009. Agrobacterium-mediated genetic transformation of pineapple var. Queen using a novel encapsulation-based antibiotic selection technique. *Plant Cell, Tissue and Organ Culture* 97:295-302.
- Ghosh, S.N. and Tarai, R.K., 2009. Influence of nitrogen levels on pineapple grown as intercrop in laterite soil. *Indian Journal of Fertilisers* 5:27-28.
- Gil, M.I. and Kader, A.A., 2008. Fresh-cut fruit and vegetables, p. 475-504, Woodhead Publishing in Food Science, Technology and Nutrition. Woodhead Publishing Ltd, Cambridge.
- Gonzaga, V. and Santos, J.M., 2009. Detection of *Pasteuria thornei* in *Pratylenchus brachyurus* and *P. zaeae*. *Nematologia Brasileira* 33:103-105.
- Gough, K.V. and Fold, N., 2009. Rise and fall of smallholder pineapple production in Ghana: changing global markets, livelihoods and settlement growth, p. 59-73. Routledge, Abingdon.
- Guadarrama, A. and Carreno, M., 2009. Ripening and post-harvest behaviour of two regional pineapple cultivars of the amazonas region in Venezuela. *Acta Horticulturae* 822:269-275.
- Gutiérrez-oliva, V.F., Abud-Archila, M., Flores-Pérez, A., Alvarez-Solis, J.D., and Gutiérrez-Miceli, F.A., 2009. Influence of arbuscular mycorrhizal fungi on the growth of pineapple vitro plantlet (*Ananas comosus* (L.) Merr) with different phosphorus levels. *Gayana Botanica* 66:1-9.
- Hamad, A.M. and Taha, R.M., 2009. Effect of explants density on the in vitro proliferation and growth of separated and cluster shoots of smooth cayenne pineapple (*Ananas comosus* L. Merr.). *Asian Journal of Plant Sciences* 8:313-317.
- He, Y. and Ou, S., 2008. Determination of nutrients in pineapple peduncle. *Modern Food Science and Technology* 24:1061-1062, 1972.
- Heenkenda, H.M.S. and Sangakkara, U.R., 2008. Plant morphological diversity of Kew and Mauritius pineapple eco-types in Sri Lanka. *Indian Journal of Horticulture* 65:393-398.
- Hu, J.S., Sether, D.M., Melzer, M.J., Subere, C.V., Cheah, K., Chen, Y., Li, Q., Borth, W., Wang, I.C., Nagai, C., and Wang, M.L., 2009. Characterization and management of pineapple mealybug wilt associated viruses. *Acta Horticulturae* 822:185-189.
- Huang, J., Bin, S., Xiang, M., Zhou, X., and Wang, L., 2008. Identification of stem root and root rot pathogens isolated from *Anthurium andraeanum* and *Ananas comosus* var. *variegata*. *Acta Phytophylacica Sinica* 35:479-480.
- Huang, K., Chiang, J., Chiu, H., Huang, Y., and Huang, C., 2009. A study on soil physical and chemical properties under different vegetation in Laopi terrace. *Crop, Environment & Bioinformatics* 6:51-60.
- Huang, W. and Zhang, Y., 2009. Abscisic acid distribution during somatic embryogenesis in pineapple callus. *Journal of the Agricultural Association of Taiwan* 10:52-62.
- Huerta, E., Kampichler, C., Geissen, V., Ochoa-Gaona, S., Jong, B.d., and Hernández-Daumás, S., 2009. Towards an ecological index for tropical soil quality based on soil macrofauna. *Pesquisa Agropecuária Brasileira* 44:1056-1062.
- Ines, M.B.C., Magdalita, P.M., Viãa, C.D.d., Cruz, F.S.J.d., and Villegas, V.N., 2009. Randomly amplified polymorphic DNA (RAPD) analysis to reveal genetic relationships among pineapple (*Ananas comosus* (L.) Merr.) genotypes. *Philippine Journal of Crop Science* 34:1-10.
- Jai, P., Bhattacharyya, S., Krishnendu, C., Roy, S., Das, S.P., and Singh, N.P., 2009. PQM-1: a newly developed superior clone of pineapple for northeastern India as evident through phenotype, fruit quality and DNA polymorphism. *Scientia Horticulturae* 120:288-291.
- Jamal, P., Tompong, M.F., and Alam, M.Z., 2009. Optimization of media composition for the production of bioprotein from pineapple skins by liquid-state bioconversion. *Journal of Applied Sciences* 9:3104-3109.
- Jaya, S. and Das, H., 2009. Glass transition and sticky point temperatures and stability/mobility diagram of fruit powders. *Food and Bioprocess Technology* 2:89-95.
- Jilani, V.A.K. and Jasmine, G.I., 2009. Preparation of chitin from shrimp shell waste using proteolytic extract of pineapple, *Ananas comosus* M. *Green Farming* 2:417-419.
- Karim, O.R., Awonorin, S.O., and Sanni, L.O., 2008. Effect of pretreatments on quality attributes of air-dehydrated pineapple slices. *Journal of Food Technology* 6:158-165.
- Kingsly, A.R.P., Balasubramaniam, V.M., and Rastogi, N.K., 2009. Effect of high-pressure processing on texture and drying behavior of pineapple. *Journal of Food Process Engineering* 32:369-381.
- Kluge, R.A., Roulet, M.C., Ongarelli, M.G., Aguila, J.S.d., and Heiffig, L.S., 2009. Physicochemical changes of pineapple submitted to different mechanical injuries. *Acta Horticulturae* 822:285-290.
- Lam Viet, D. and JenÁ-Á?ek, V., 2008. The study on penetration capacity of pineapple products into USA's market. *Agricultura Tropica et Subtropica* 41:12-16.
- Leal, A.J.F., Hora, R.C.d., Tonin, T.A., and Boliani, A.C., 2009. Economic feasibility of pineapple culture in the Caiuá sandstone area, northwestern Paraná State. *Acta Scientiarum - Agronomy* 31:353-358.

- Leão, A.L., Machado, I.S., Souza, S.F.d., and Soriano, L., 2009. Production of curaua (*Ananas erectifolius* L.B. Smith) fibers for industrial applications: characterization and micropropagation. *Acta Horticulturae* 822:227-238.
- Lebeau, F., Imele, J.P., Teisson, C., and Delhove, G., 2009. Efficiency of the *Ananas comosus* (L.) Merr. flower induction treatment based on ethylene enriched activated carbon (TIFBio). *Biotechnologie, Agronomie, Soci  t   et Environnement* 13:395-400.
- Liang, X., Chen, X., Kheng, C., Diane, M.S., Qi, L., and John, H., 2009. Establishment of direct somatic embryogenesis system through transverse thin cell layer culture in pineapple (*Ananas comosus* L. Merrill). *Acta Horticulturae Sinica* 36:1597-1602.
- Lin, C.H., Maruthasalam, S., Shiu, L.Y., Lien, W.C., Loganathan, M., Yu, C.W., Hung, S.H., Ko, Y., and Chen, Y.Y., 2009. Physical and chemical manipulation of flowering in pineapple. *Acta Horticulturae* 822:117-123.
- Liu, C., Liu, Y., Yi, G., Liao, M., Wei, F., and Zhu, S., 2009. Physiological and biochemical mechanisms of different organic fertilizer applications affecting pineapple plant growth. *Acta Botanica Boreali-Occidentalia Sinica* 29:2527-2534.
- Liu, C., Liu, Y., Yi, G., Zhang, G., Zhong, Y., Jiang, B., and Luo, M., 2009. Comparison of aroma components in pineapple fruits ripened in summer and autumn. *Acta Botanica Boreali-Occidentalia Sinica* 29:397-401.
- Liu, D., Wei, J., Wang, Y., and Li, S., 2008. The wild and cultivated fruit crop resources in Hainan province. *South China Fruits*:34-37.
- Lopez Herrera, M., WingChing-Jones, R., and Rojas-Bourrill  n, A., 2009. Fermentative and nutritional characteristics of pineapple plant straw silage. *Agronomia Costarricense* 33:1-15.
- Machado, F.L.C., Alves, R.E., Figueiredo, R.W., and Teixeira, A.S., 2009. Quality maintenance of ripe pineapple as affected by application of wax associated to 1-methylcyclopropene. *Acta Horticulturae* 822:261-268.
- Maia, V.M., Almeida J  nior, A.B., Mizobutsi, G.P., Mizobutsi, E.H., and Pacheco, D.D., 2009. Fruit and planting material production by irrigated 'Perola' pineapple in response to planting spacing under semi-arid conditions. *Acta Horticulturae* 822:125-130.
- Mainoo, N.O.K., Barrington, S., Whalen, J.K., and Sampedro, L., 2009. Pilot-scale vermicomposting of pineapple wastes with earthworms native to Accra, Ghana. *Bioresourcetechnology* 100:5872-5875.
- Maliwan, P., 2009. Effects of pineapple bran in broiler rations during 0-3 week, p. 141-147. Kasetsart University, Bangkok.
- Mangara, A., Adopo, A.A.N., Kouame, N.M.T., and Kehe, M., 2009. Effect of polythene and cover crop *Mucuna pruriens* (L.) DC. in the control of weeds in pineapple (*Ananas comosus* (L.) Merr.) in C  te d'Ivoire. *Journal of Applied Biosciences* 22:1326-1332.
- Mangara, A., Adopo, A.N., Boraud, M.K., Kobenan, K., Lejoly, J., and Traore, D., 2008. Inventory of the weedy flora of pineapple plantations (*Ananas comosus* (L.) Merr.) located around the city of Bonoua in lowland C  te d'Ivoire. *Agronomie Africaine* 20:23-35.
- Marques, L.G., Prado, M.M., and Freire, J.T., 2009. Rehydration characteristics of freeze-dried tropical fruits. *LWT - Food Science and Technology* 42:1232-1237.
- Matos, A.P.d. and Reinhardt, D.H., 2009. Pineapple in Brazil: characteristics, research and perspectives. *Acta Horticulturae* 822:25-36.
- Matos, A.P.d., Sanches, N.F., Souza, L.F.d.S., Elias J  nior, J., Teixeira, F.A., and Siebeneichler, S.C., 2009. Cover crops on weed management in integrated pineapple production plantings. *Acta Horticulturae* 822:155-160.
- Matos, A.P.d., Sanches, N.F., Teixeira, F.A., and Elias J  nior, J., 2009. Integrated management of fusariosis in pineapple fields under integrated production system. *Acta Horticulturae* 822:199-204.
- Melo, E.d.A., Maciel, M.I.S., Lima, V.L.A.G.d., and Ara  jo, C.R.d., 2008. Total phenolic contents and antioxidant capacity of the frozen fruit pulps. *Alimentos e Nutri  o* 19:67-72.
- Mezadri, T., Bramorski, A., Faria, M.P.d., Cunha, T.I.d., and Costa, A.d.A.d.S.d., 2008. Total polyphenol index and vitamin C concentration in fruit pulps produced in Santa Catarina, Brazil. *Higiene Alimentar* 22:22-25.
- Minal, M., Nagi, L., and Ganapathi, T.R., 2009. Agrobacterium-mediated transformation of pineapple (*Ananas comosus* L. Merr.) leaf bases with MSI-99, a magainin analogue. *International Journal of Fruit Science* 9:106-114.
- Moda, E.M., Pilon, L., Zocchi, S.S., and Spoto, M.H.F., 2008. Physical-chemical and sensorial quality of irradiated and minimally processed pineapple. *Boletim do Centro de Pesquisa e Processamento de Alimentos* 26:267-276.
- Mohamadu Boyie, J., Wan Sulaiman, W.H., Jamal, T., Mohd Fauzi, R., Rajan, A., Sung, T., and Osumanu Haruna, A., 2009. A simulation model estimates of the intercropping advantage of an immature-rubber, banana and pineapple system. *American Journal of Agricultural and Biological Sciences* 4:249-254.
- Mohammad, A., Rajeev, B., and Karim, A.A., 2009. UV radiation-induced changes of antioxidant capacity of fresh-cut tropical fruits. *Innovative Food Science & Emerging Technologies* 10:512-516.
- Mondino, E.A., Tavares, O.C.H., Ebeling, A.G., Figueira, A.F., Quintero, E.I., and Berbara, R.L.L., 2009. Evaluation of nematodes communities in organic agricultural system. *Acta Scientiarum - Agronomy* 31:509-515.
- Moryadee, A. and Pathom-Aree, W., 2008. Isolation of thermotolerant acetic acid bacteria from fruits for vinegar production. *Research Journal of Microbiology* 3:209-212.
- Munawar, S.S., Umemura, K., and Kawai, S., 2008. Manufacture of oriented board using mild steam treatment of plant fiber bundles. *Journal of Wood Science* 54:369-376.
- Murray, D.N.A. and Manicom, B., 2009. Development of an organic pineapple cultivation strategy for the Eastern Cape Region of South Africa. *Acta Horticulturae* 822:107-116.
- Nath, J.C., Saud, B.K., Chowdhury, D., Deka, K.K., and Sarma, U.J., 2008. Coconut based high density multispecies cropping system in Assam. *Journal of Plantation Crops* 36:98-102.
- Neto, J.F.d.B., Pereira, W.E., S  bri  nho, R.G.d., Barbosa, J.A., Costa, D.d.S., Lacerda, J.S., Santos, D.P.d., and Vieira, D.d.O., 2009. Commercialization forms and organization of pineapple producers in the State of Para  ba, Brazil. *Acta Horticulturae* 822:313-316.
- Neuteboom, L.W., Matsumoto, K.O., and Christopher, D.A., 2009. An extended AE-rich N-terminal trunk in secreted pineapple cystatin enhances inhibition of fruit bromelain and is posttranslationally removed during ripening. *Plant Physiology* 151:515-527.

- Nguyen Thi Hong, N., Nguyen Van, H., and Preston, T.R., 2009. Ensiling with or without additives to preserve pineapple residue and reduce pollution of the environment. *Livestock Research for Rural Development* 21:96.
- Nilprapruck, P., Pradisthakarn, N., Authanithe, F., and Keebjan, P., 2008. Effect of exogenous methyl jasmonate on chilling injury and quality of pineapple (*Ananas comosus* L.) cv. Pattavia. *Silpakorn University Science and Technology Journal* 2:33-42.
- Oliveira, M.D.d.M. and Nascimento, L.C.d., 2009. Activity evaluation of abiotic resistance inducers, chemical fungicide and natural plant extracts on black rot of pineapple, cv. pãrola. *Revista Brasileira de Fruticultura* 31:84-89.
- Pedreira, A.C.d.C., Naves, R.V., and Nascimento, J.L.d., 2008. Seasonal variation of the pineapple cv. Perola quality in Goiania, Goias State, Brazil. *Pesquisa Agropecuária Tropical* 38:262-268.
- Pegna, F.G. and Fuentes, J.S.P.d.C., 2009. Development of a vertical axis shredder for pineapple field clearing. *Acta Horticulturae* 822:177-182.
- Peng, Q., Wang, L., and Pan, S., 2009. Study of the brewing process of the pineapple wine. *Modern Food Science and Technology* 25:687-689, 395.
- Pereira, E.S., Regadas Filho, J.G.L., Arruda, A.M.V.d., Mizubuti, I.Y., Villarreal, A.B.S., Pimentel, P.G., and Cândido, M.J.D., 2008. NRC (2001) equations to predict the energy value of agroindustry byproducts in Brazilian northeast. *Revista Brasileira de Saude e Producao Animal* 9:258-269.
- Pereira, E.S., Regadas Filho, J.G.L., Freitas, E.R., Neiva, J.N.M., and Cândido, M.J.D., 2009. Energetic value from by-product of the Brazil agroindustria. *Archivos de Zootecnia* 58:455-458.
- Pereira, M.A.B., Siebeneichler, S.C., Lorenãoni, R., Adorian, G.C., Silva, J.C.d., Garcia, R.B.M., Pequeno, D.N.L., Souza, C.M.d., and Brito Filho, R.F.d., 2009. Quality of pineapple commercialized by Cooperfruto - Miranorte/Tocantins/Brazil. *Revista Brasileira de Fruticultura* 31:1048-1053.
- Poelman, A., Mojet, J., Lyon, D., and Sefa-Dedeh, S., 2008. The influence of information about organic production and fair trade on preferences for and perception of pineapple. *Food Quality and Preference* 19:114-121.
- Quaggio, J.A., Teixeira, L.A.J., Cantarella, H., Mellis, E.V., and Sigrist, J.M., 2009. Post-harvest behaviour of pineapple affected by sources and rates of potassium. *Acta Horticulturae* 822:277-284.
- Rabie, E.C. and Tustin, H.A., 2009. The effect of different cover crops on nematode populations and yield in 'Queen' pineapple cultivation. *Acta Horticulturae* 822:205-213.
- Ramadan-Hassanien, M.F., 2008. Total antioxidant potential of juices, beverages and hot drinks consumed in Egypt screened by DPPH in vitro assay. *Grasas y Aceites (Sevilla)* 59:254-259.
- Ramirez, A. and Pacheco de Delahaye, E., 2009. Functional properties of starches with high dietetic fiber content obtained from pineapple, guava and sourp. *Interciencia* 34:293-298.
- Ramos, A.M., Quintero, A.C.F., Faraoni, A.S., Soares, N.d.F.F., and Pereira, J.A.M., 2008. Effect of packing type and storage time on the physical-chemical and microbiological quality of dehydrated pineapple. *Alimentos e Nutriãõ* 19:259-269.
- Ramos, M.J.M., Monnerat, P.H., Carvalho, A.J.C.d., Pinto, J.L.d.A., and Silva, J.A.d., 2009. Visual symptoms of macronutrients and boron deficiency in 'Imperial' pineapple. *Revista Brasileira de Fruticultura* 31:252-256.
- Ramos, M.J.M., Monnerat, P.H., Pinho, L.G.d.R., Carvalho, A.J.C., and Silva, J.A.d., 2009. Morphological characteristics of 'Imperial' pineapple fruits under deficiency of macronutrients and boron. *Acta Horticulturae* 822:147-153.
- Ramos, M.J.M., Monnerat, P.H., Pinho, L.G.d.R., and Pinto, J.A.d.A., 2009. Growth and flowering of 'Imperial' pineapple plants under macronutrient and boron deficiency. *Acta Horticulturae* 822:139-146.
- Reinhardt, A. and Rodriguez, L.V., 2009. Industrial processing of pineapple - trends and perspectives. *Acta Horticulturae* 822:323-328.
- Reinhardt, D.H.R.C., 2009. Proceedings of the Sixth International Pineapple Symposium, Joao Pessoa, Brazil, 18-23 November 2007. *Acta Horticulturae* 822:328 pp.
- Reungmanee-paitoon, S. and Yai-eiam, S., 2008. Production of instant beverage of the pineapple and sweetpotato. 387-389.
- Rodrigues, E., Silva, J.N.d., Souza, C.M.A.d., Lacerda Filho, A.F.d., and Chaves, M.A., 2008. Simulation of fixed-bed drying of pineapple slices. *Engenharia na Agricultura* 16:381-393.
- Rodrigues e Silva, A.F. and Zambiasi, R.C., 2008. Acceptability of conventional and light pineapple jellies obtained from agroindustrial residues. *Boletim do Centro de Pesquisa e Processamento de Alimentos* 26:1-8.
- Rosnah, S., Wan Daud, W.R., Mohd Sobri, T., and Osman, H., 2009. Physico-mechanical properties of the Jospine pineapple fruits. *Pertanika Journal of Science & Technology* 17:117-123.
- Sanewski, G.M., 2009. Breeding Ananas for the cut-flower and garden markets. *Acta Horticulturae* 822:71-78.
- Sanewski, G.M., 2009. The effect of different levels of inbreeding on self-incompatibility and inbreeding depression in pineapple. *Acta Horticulturae* 822:63-70.
- Sangjindavong, M., Mookdasanit, J., Wilaipun, P., Chuapoehuk, P., and Akkanvanitch, C., 2009. Using pineapple to produce fish sauce from Surimi waste. *Kasetsart Journal, Natural Sciences* 43:791-795.
- Santos, P.H.S. and Silva, M.A., 2009. Kinetics of L-ascorbic acid degradation in pineapple drying under ethanolic atmosphere. *Drying Technology* 27:947-954.
- Sanusi, R.A., Ogunro, Y., and Nwozoh, S., 2008. Effect of storage time on ascorbic acid content of some selected "Made in Nigeria" fruit preserves. *Pakistan Journal of Nutrition* 7:730-732.
- Senhor, R.F., Souza, P.A.d., Carvalho, J.N.d., Silval, F.L.d., and Silva, M., 2009. Factors for pre and post-harvest affecting the fruits and vegetables in postharvest. *Revista Verde de Agroecologia e Desenvolvimento Sustentável* 4:13-21.
- Silva, A.P.d., Alvarez, V.V.H., Souza, A.P.d., Neves, J.C.L., Novais, R.F., and Dantas, J.P., 2009. Fertilizer and lime recommendation system for pineapple - fertcalc-abacaxi. *Revista Brasileira de Ciãncia do Solo* 33:1269-1280.
- Silva, L.L.d., Costa, R.F.d., Campos, J.H.B.d.C., and Dantas, R.T., 2009. Influence of precipitations on agricultural productivity in Paraãba State. *Revista Brasileira de Engenharia Agrãcola e Ambiental* 13:454-461.

- Silveira, E., Souza JÃnior, M.E., Santana, J.C.C., Chaves, A.C., Porto, A.L.F., and Tambourgi, E.B., 2009. Expanded bed adsorption of bromelain (E.C. 3.4.22.33) from *Ananas comosus* crude extract. *Brazilian Journal of Chemical Engineering* 26:149-157.
- Singh, D.B., 2009. Comparative study of organic and conventional farming of pineapple. *Asian Journal of Horticulture* 4:1-4.
- Sirousazar, M., Mohammadi-Doust, A., and Achachlouei, B.F., 2009. Mathematical investigation of the effects of slicing on the osmotic dehydration of sphere and cylinder shaped fruits. *Czech Journal of Food Sciences* 27:95-101.
- Slongo, A.P. and AragÃo, G.M.F.d., 2008. Behaviour of *Neosartorya fischeri* ascospores in pineapple juice. *Boletim do Centro de Pesquisa e Processamento de Alimentos* 26:51-60.
- Soares, S.A.G., Mariano, R.d.L.R., Cavalcante, U.M.T., and Maia, L.C., 2009. Effect of bacteria on germination of arbuscular mycorrhizal fungi and their co-inoculation on pineapple seedlings. *Caatinga* 22:31-38.
- Soler, A., Marie-Alphonsine, P.A., Topart, P., and Queneherve, P., 2009. How to evaluate the resistance or tolerance of pineapple varieties to nematodes? *Acta Horticulturae* 822:191-198.
- Sossou, S.K., Ameyapoh, Y., Karou, S.D., and Souza, C.d., 2009. Study of pineapple peelings processing into vinegar by biotechnology. *Pakistan Journal of Biological Sciences* 12:859-865.
- Souza, F.V.D., Cabral, J.R.d.S., Souza, E.H.d., Silva, M.d.J., Santos, O.S.N., and Ferreira, F.R., 2009. Evaluation of F1 hybrids between *Ananas comosus* var. *ananassoides* and *Ananas comosus* var. *erectifolius*. *Acta Horticulturae* 822:79-84.
- Souza, O.P.d., Teodoro, R.E.F., Melo, B.d., and Torres, J.L.R., 2009. Fruit quality and productivity of pineapple under different planting densities and irrigation levels. *Pesquisa Agropecuária Brasileira* 44:471-477.
- Sriporaya, S., 2009. Pineapple hybridization and selection in Thailand. *Acta Horticulturae* 822:57-62.
- Sriwichailamphan, T., Sriboonchitta, S., Wiboonpongse, A., and Chaovanapoonphol, Y., 2008. Factors affecting good agricultural practice in pineapple farming in Thailand. 325-334.
- Sudhanshu, S., Mishra, B.B., Ramesh, C., and Arun, S., 2009. Shelf stable intermediate moisture pineapple (*Ananas comosus*) slices using hurdle technology. *LWT - Food Science and Technology* 42:1681-1687.
- Taylor, R., 2009. Harvestable hedgerows encourage erosion change. *Partners in Research for Development*:12-15.
- Teixeira, L.A.J., Quaggio, J.A., and Zambrosi, F.C.B., 2009. Preliminary DRIS norms for 'Smooth Cayenne' pineapple and derivation of critical levels of leaf nutrient concentrations. *Acta Horticulturae* 822:131-138.
- Thakor, N.J. and Sawant, A.A., 2008. Effect of sucrose concentration and temperature on osmotic dehydration of pineapple slices. *Agriculture Update* 3:417-420.
- Uckiah, A., Goburdhun, D., and Ruggoo, A., 2009. Vitamin C content during processing and storage of pineapple. *Nutrition & Food Science* 39:398-412.
- Vagneron, I., Faure, G., and Loeillet, D., 2009. Is there a pilot in the chain? Identifying the key drivers of change in the fresh pineapple sector. *Food Policy* 34:437-446.
- Ventura, J.A., Costa, H., Cabral, J.R.S., and Matos, A.P.d., 2009. 'Vitoria': new pineapple cultivar resistant to fusariosis. *Acta Horticulturae* 822:51-55.
- Viana, E.S., Amorim, T.S., Oliveira, L.A., Reinhardt, D.H., Silveira, F.G.F., Santana, F.A., and Santana, P.N., 2009. Physicochemical characteristics of minimally processed 'Perola' pineapples treated with edible coatings. *Acta Horticulturae* 822:291-297.
- Villalobos Moreno, A., Agudelo M, J.C., Barajas S, A.F., Herrera Q, S., and CÃspedes, J.C., 2009. Morphological changes throughout the ontogeny of the pupal stage of *Melanomoma viatrix* (Diptera: Richardiidae). *Revista Colombiana de Entomología* 35:98-100.
- Vinodhini, S. and Malathy, N.S., 2009. Bioprospecting of plants fibre of Coimbatore district of TamilNadu. *International Journal of Plant Sciences (Muzaffarnagar)* 4:444-445.
- Wang, M.-L., Uruu, G., Xiong, L., He, X., Nagai, C., Cheah, K., Hu, J., Nan, G.-L., Sipes, B., Atkinson, H., Moore, P., Rohrbach, K., and Paull, R., 2009. Production of transgenic pineapple (*Ananas cosmos* (L.) Merr.) plants via adventitious bud regeneration *In Vitro Cellular and Developmental Biology - Plant* 45:112-121.
- Waughon, T.G.M. and Pena, R.d.S., 2008. Thin layer drying model of pineapple's fibrous residue. *Boletim do Centro de Pesquisa e Processamento de Alimentos* 26:297-306.
- Wei, C., Chen, J., Liu, S., Sun, G., and Zang, X., 2009. Primary research of effect of foliar spray of nutrition on pineapple aroma. *Southwest China Journal of Agricultural Sciences* 22:382-384.
- Wen, S., Cheng, Y., Du, B., Liang, J., and Yang, G., 2008. Effects of the sterilization methods on the physico-chemical properties of pineapple juice. *Modern Food Science and Technology* 24:977-980.
- Xaud, M.R., Xaud, H.A.M., MourÃo, M., Jr., Bendahan, A.B., and Carvalho, A.L.M., 2009. Evaluation of pineapple crop used as firebreak in Roraima, Amazonia. *Acta Horticulturae* 822:245-249.
- Xavier, V.L.S.M., Oliveira Junior, L.F.G.d., Berger, R.M., Bispo, W.M.d.S., Stein, Ã.M., Santos, R.B.d., and Matsumoto, S.T., 2009. Assessment of the potential of sanitizing agents for the minimal processing of 'Perola' pineapple. *Revista Brasileira de Armazenamento* 34:26-32.
- Yabor, L., Valle, B., Carvajal, C., Aragón, C., Hernández, M., González, J., Daquinta, M., Arencibia, A., and Carlos Lorenzo, J., 2009. Characterization of a field-grown transgenic pineapple clone containing the genes chitinase, AP24, and bar. *In Vitro Cellular & Developmental Biology - Plant* 46:1-7.
- Yang, D., Zhang, T., Ma, Q., Duan, J., and Liu, C., 2009. Study of *Agaricus bisporus* tyrosinase inhibitors extracted from pineapple and their anti-browning effects. *Modern Food Science and Technology* 25:1032-1034.
- Yang, X., Cai, Y., Wu, Q., and Sun, G., 2009. Cloning and expression analysis of a zinc finger protein gene AcRCHY1 from pineapple. *Acta Horticulturae Sinica* 36:1589-1596.
- Yoshimoto, M., Nishiba, Y., Kurata, R., Ikemiya, H., Nakasone, F., Shoda, M., Takeuchi, M., Higa, A., and Uehara, E., 2008. Physiological functions of pineapple-peel vinegar. *Report of the Kyushu Branch of the Crop Science Society of Japan*:62-64.

- Youryon, P., Wongs-Aree, C., McGlasson, W.B., Glahan, S., and Kanlayanarat, S., 2008. Internal browning occurrences of 'Queen' pineapple under various low temperatures. *Acta Horticulturae* 804:555-560.
- Zhang, H., Wu, Q., Sun, W., Zan, F., and Dou, M., 2009. Method of modified CTAB for DNA extraction from pineapple. *Southwest China Journal of Agricultural Sciences* 22:440-443.
- Zhang, N., ZeTan, Zhang, X., and Ren, M., 2009. The pineapple diseases and pests in China and their control. *South China Fruits*:52-55.
- Zhang, X., Du, L., Sun, G., Wei, C., Liu, S., and Xie, J., 2009. Changes of aroma components in Yellow Mauritius pineapple during fruit development. *Journal of Fruit Science* 26:245-249.

Pineapple Reference Database

A pineapple references database containing over 7,600 references on pineapple is maintained by the editor. Literature searches of the database on specific topics, including abstracts where available, can be obtained by contacting Duane Bartholomew at duaneb@hawaii.edu. ♦

Instructions to Contributors to Pineapple News

All contributions should be written in English. Editing assistance will be provided on request.

Preferred contributions include:

- Timely news about research on issues related to culture, processing, storage, and marketing of pineapple.
- New, interesting, or unique problems encountered by growers.
- Country or status reports on the local pineapple industry.
- If uncertain about the suitability of material for the newsletter, contact the editor.

If possible, please send contributions by E-mail as attached files in MS Word or rich text format or on floppy disks. When sending printed copy, be sure that it is clean and sharp so it can be scanned to speed conversion to a wordprocessor format.

Article length: Papers usually should be no longer than 4 double-spaced pages in 12 point font or equivalent, not including tables, figures and photos. If longer than 4 pages, please contact the editor. There is no limit on the number of articles that can be submitted. However, acceptance and publication is at the discretion of the editor.

Tables: The preferred table format is columns separated by tabs. Authors may be asked to revise tables not in the requested format.

Photographs: Submit photographs that can be scanned or provide digital files in jpeg format with a minimum resolution of 300 dpi so they can be printed with acceptable resolution in grey scale with a laser printer.

Mail contributions and inquiries to: **D.P. Bartholomew, Dept. of NREM, Univ. of Hawaii, 1910 East-West Rd., Honolulu, HI 96822 U.S.A.** (Phone (808) 956-7568; Fax (808) 956-6539; E-mail: duaneb@hawaii.edu.)

Pineapple News is available on the Web at: <http://tpss.hawaii.edu/pineapple/pineappl.htm>. ♦

Pineapple News is published by the University of Hawaii, College of Tropical Agriculture and Human Resources, Dept. of Tropical Plant and Soil Science. The contributions of Valent Biosciences and by various individuals to the University of Hawaii Foundation help support the costs associated with the publication of the newsletter. Reference to commercial products and services is made for the convenience of readers with the understanding that no discrimination is intended and no endorsement by the University of Hawaii and their employees is implied.

Information in this newsletter is public property and may be reprinted without permission.

The University of Hawaii, College of Tropical Agriculture and Human Resources is an Equal Opportunity and Affirmative Action Employer.