

NEWS & VIEWS

A regional newsletter published by the
Potash & Phosphate Institute (PPI) and the
Potash & Phosphate Institute of Canada (PPIC)



Dr. T.W. Bruulsema,
Eastern Canada and
Northeast U.S. Director
May 1998

Mixed Hay Fields Respond to NPK in Fertilizer and Manure

ACROSS Eastern Canada and the northeastern U.S., more than 5 million acres of land produces mixed hay from forage crops other than alfalfa (Table 1). Estimates of the economic value of the crop vary across the region, but average about US\$60 per ton. The hay crop is often undervalued. Therefore, fertility needs receive little attention on much of this land. The lost opportunities arising from under-fertilizing could be in the neighborhood of \$300 million, assuming that yields would improve by 50 percent if nutrient inputs were increased.

Forages such as timothy, orchardgrass and reed canary-grass respond to nitrogen (N), phosphorus (P), and potassium (K). Their productivity can exceed 4 tons/A of hay, but average yields are much lower (Table 1). Yields of 4 tons/A or more can quickly deplete available nutrients from the soil. A typical crop of mixed hay removes about 13 lb of P_2O_5 and 50 lb of K_2O per ton of dry matter harvested.

Table 1. Estimated hay production, other than alfalfa, in eastern Canada and northeastern U.S.A. (1995-1996).

	Area, acres	Production, tons	Value, \$US	Yield, tons/A
Ontario	1,275,000	2,780,000	129,000,000	2.18
Quebec	1,075,000	2,365,000	110,000,000	2.20
Atlantic	225,000	439,000	22,000,000	1.95
New England	566,000	1,069,000	99,000,000	1.89
New York	870,000	1,740,000	118,000,000	2.00
Pennsylvania	1,130,000	1,674,000	135,000,000	1.48
New Jersey	120,000	200,000	17,000,000	1.66
Total	5,261,000	10,267,000	630,000,000	1.95

Recently in Maine, research compared manure and fertilizer sources of NPK for their impact on yield from a mixed hay field. Manure provided 125 lb/A of N, to supply the first and second growth of the sward. It also supplied about 95 lb/A of P_2O_5 and 250 lb/A of K_2O . A normal recommendation for the soil would have been 30 lb/A of P_2O_5 and 100 lb/A of K_2O . Fertilizer was applied matching the available NPK rate of the manure, and, in another treatment, matching only the available N. Applications were split so that 60 percent was applied in mid-May and 40 percent after the first cut in mid-June. Additionally, all treatments received 50 lb/A of N as ammonium nitrate (NH_4NO_3) following the second cut.

The fertilizer source produced the greatest yield response (Figure 1). Most of the yield response was due to N alone, but added P and K produced an additional yield increase that was statistically significant when the first two years were considered. While the yield response to added P and K was not large enough to justify the fertilizer cost in these first two years, in the longer term the response would be larger. The crop removed more P and K than would have been supplied in a normal recommendation. The N only treatment caused a sharp reduction in soil test levels for both P and K. For sustainable yield in the longer term, the use of P and K would provide economic returns.

Manure alone also produced a yield response. Hay yield when manured was 88 percent of that when fertilized with NPK, and 91 percent of that fertilized only with N. The fact that manured fields

Co-authors:

Dr. Tom W. Bruulsema
Eastern Canada and
Northeast U.S. Director, PPI
18 Maplewood Drive
Guelph, Ontario, Canada N1G 1L8
Phone: (519) 821-5519
E-mail: bruulsem@ppi-far.com

Dr. Timothy S. Griffin,
Sustainable Agriculture & Forage Crops
University of Maine, Cooperative Extension
495 College Avenue
Orono, Maine 04473-1294
Phone: (207) 581-2942
Fax: (207) 581-1301

Dr. Mary Wiedenhoft
Assoc. Professor of Agronomy
University of Maine
5722 Deering Hall
Orono, Maine 04469-5722
Phone: (207) 581-2951
Fax: (207) 581-2999

Eric Giberson
Aroostook Farm
59 Houlton Road
Presque Isle, Maine 04769
Phone: (207) 764-3122
Egiber51@maine.maine.edu

