

Table 14. Use of lures and repellents in vineyards for the management of MALB, 8/31/09 to 9/17/09.

| Location / Treatment | <i>n</i> | MALB per cluster | Proportion of clusters with fresh damage | MALB per yellow stick card |
|------------------------------|----------|------------------|--|----------------------------|
| <i>Hastings, MN 1</i> | | | | |
| Lure/trap/repellent area | 6 | 0.36 | 0.63 | 4.25 |
| Untreated check area | 6 | 0.49 | 0.66 | -- |
| <i>Hastings, MN 2</i> | | | | |
| Lure/trap/repellent area | 6 | 0.31 | 0.74 | 5.12 |
| Untreated check area | 6 | 0.41 | 0.77 | -- |
| <i>Stillwater, MN</i> | | | | |
| Lure/trap/repellent area | 6 | 0.00 | 0.26 | 0.70 |
| Untreated check area | 6 | 0.00 | 0.25 | -- |
| <i>Somerset, WI 1</i> | | | | |
| Lure/trap/repellent area | 6 | 0.00 | 0.04 | 0.88 |
| Untreated check area | 6 | 0.00 | 0.01 | -- |
| <i>Somerset, WI 2</i> | | | | |
| Lure/trap/repellent area | 6 | 0.00 | 0.06 | 0.36 |
| Untreated check area | 6 | 0.00 | 0.05 | -- |
| <i>Total</i> | | | | |
| Lure/trap/repellent area | 30 | 0.13 | 0.33 | 2.14 |
| Untreated check area | 30 | 0.17 | 0.33 | -- |
| | | NS | NS | NS |

NS = not significant.

Table 15. Use of lures and repellents in vineyards for the management of MALB, 8/26/10 to 9/9/10.

| Location / Treatment | <i>n</i> | MALB per cluster | Proportion of clusters with fresh damage | MALB per yellow stick card |
|---------------------------------|----------|------------------|--|----------------------------|
| <i>Hastings, MN 1</i> | | | | |
| Repellent area | 5 | 0.01 | 0.50 | -- |
| Between lure and repellent area | 5 | 0.00 | 0.47 | 0.47 |
| Untreated check area | 5 | 0.00 | 0.50 | -- |
| <i>Hastings, MN 2</i> | | | | |
| Repellent area | 5 | 0.01 | 0.90 | -- |
| Between lure and repellent area | 5 | 0.04 | 0.86 | 1.05 |
| Untreated check area | 5 | 0.03 | 0.85 | -- |
| <i>Stillwater, MN</i> | | | | |
| Lure/trap | 5 | 0.01 | 0.61 | 0.03 |
| Untreated check | 5 | 0.00 | 0.51 | -- |
| <i>Somerset, WI</i> | | | | |
| Lure/trap | 5 | 0.00 | 0.22 | 0.03 |
| Untreated check | 5 | 0.00 | 0.30 | -- |
| | | NS | NS | NS |

NS = not significant.

Table 16. Use of lures and repellents in vineyards for the management of MALB, 8/30/11 to 9/9/11.

| Location / Treatment | <i>n</i> | MALB per cluster | Proportion of clusters with fresh damage | MALB per yellow stick card |
|---------------------------------|----------|------------------|--|----------------------------|
| <i>Hastings, MN</i> | | | | |
| Repellent area | 4 | 0.00 | 0.97 | -- |
| Between lure and repellent area | 4 | 0.00 | 0.84 | 0.30 |
| Untreated check area | 4 | 0.00 | 0.90 | -- |
| | | NS | NS | |
| <i>Stillwater, MN</i> | | | | |
| Lure/trap | 4 | 0.00 | 0.17 | 0.05 |
| Untreated check | 4 | 0.00 | 0.12 | -- |
| | | NS | NS | |
| <i>Somerset, WI</i> | | | | |
| Lure/trap | 4 | 0.00 | 0.09 | 0.09 |
| Untreated check | 4 | 0.00 | 0.05 | -- |
| | | NS | NS | |

Means within columns followed by the same letter are not significantly different, t-test; NS = not significant.