

Table 4. Subset of NESARE forage and feed quality analyses for the FORAGE rotation in A) 2010 and B) 2012. The main management comparison in this rotation is broadcasting (BM) vs. injecting (IM) manure. Standard errors (SE) are presented. See 2011 Annual Report for that year's data.

A) 2010		% Crude Protein			% Neutral Detergent Fiber			Net Energy of Lactation (Mcal/lb)		
Crop Entry	Harvest Date	IM	BM	(SE)	IM	BM	(SE)	IM	BM	(SE)
Corn Silage (after alf+grass)		8.20	8.10	0.16	32.50	33.77	1.34	0.84	0.83	0.01
Corn Silage (after red clov.)		8.10	8.20	0.16	30.73	31.60	1.34	0.85	0.83	0.01
Canola ^A										
Red Clover (fall cut)										
Alf+Grass Yr. 1 (cut #1)/silage ~	6/29/2010	18.57	18.13	0.84	45.00	41.13	2.16	0.62	0.66	0.02
Alf+Grass Yr. 1 (cut #2)/silage	8/3/2010	23.90	24.60	0.84	51.37	47.70	2.16	0.49	0.51	0.02
Alf+Grass Yr. 1 (cut #3)	9/14/2010	22.87	22.67	0.84	44.37	48.40	2.16	0.63	0.61	0.02

a,b: Different lowercase letters between IM and BM for a particular variable indicate a statistical difference between main management treatments at the 0.05 level.

^AFor canola, one composite sample of meal was taken for all treatments in the first year.

~ For alfalfa + grass, subsamples were taken for one crop entry because all crop entry points were new seedings.

B) 2012		% Crude Protein			% Neutral Detergent			Net Energy of Lactation		
Crop Entry	Harvest Date	IM	BM	(SE)	IM	BM	(SE)	IM	BM	(SE)
Corn Silage (after alf+grass)	9/17/2012	7.83	7.77	0.13	38.30	34.63	1.10	0.78	0.81	0.01
Cutting Alf+Grass Before Corn Silage*	5/11/2012	22.90		-	48.90		-	0.63		-
Corn Silage (after red clov.)	9/7/2012	8.37	7.53	0.29	35.90	35.73	2.19	0.81	0.81	0.02
Canola ^A	7/3/2012	40.2	37.9	-	31.3	91.8	-	0.86	0.89	-
Red Clover (fall cut)*	8/22/2012	22.0	21.3	-	49.6	51.8	-	0.53	0.50	-
SS Grass Yr. 1 (cut #1)/silage~	7/24/2012	15.15		0.75	67.05		0.95	0.45		0.01
SS Grass Yr. 1 (cut #2)/silage	8/30/2012	13.30	12.73	0.49	58.10	57.67	0.59	0.59	0.58	0.01
Alf+Grass Yr. 2 (cut #1)/silage	5/11/2012	22.90	21.35	IM: 1.15 BM: 0.81	41.30	41.95	IM: 3.56 BM: 2.52	0.69	0.68	IM: 0.03 BM: 0.02
Alf+Grass Yr. 2 (cut #2)/silage	6/8/2012	20.90	20.35	IM: 1.15 BM: 0.81	48.20	46.80	IM: 3.56 BM: 2.52	0.64	0.65	IM: 0.03 BM: 0.02
Alf+Grass Yr. 2 (cut #3)/hay	7/9/2012	16.70	17.45	IM: 1.15 BM: 0.81	55.40	54.00	IM: 3.56 BM: 2.52	0.58	0.58	IM: 0.03 BM: 0.02
Alf+Grass Yr. 2 (cut #4)/silage	8/3/2012	25.73	21.97	0.66	44.57	54.00	2.06	0.63	0.55	0.02
Alf+Grass Yr. 2 (cut #5)/hay	9/6/2012	21.10	20.15	IM: 1.15 BM: 0.81	46.30	51.75	IM: 3.56 BM: 2.52	0.58	0.56	IM: 0.03 BM: 0.02

a,b: Different lowercase letters between IM and BM for a particular variable indicate a statistical difference between main management treatments at the 0.05 level.

^AFor canola, results shown are the averages from three (IM) or two (BM) representative sample(s) and were not analyzed statistically.

~Only 1-2 samples were taken, so it was not analyzed with SAS.

~Only 1 sample per main mgt trmt was taken; composited mean and SE were generated in Excel.