

Penn State Extension

Can we feed less protein to dairy cows, save money, and help the environment?

extension.psu.edu

The Pennsylvania State University Ammonia Mitigation Project

In 2009, the Northeast Sustainable Agriculture Research and Education (NE SARE) program funded a 3-year project targeting reduction of ammonia emissions from dairy farms in Pennsylvania.



Feeding protein to dairy cows in excess of their requirements is no longer an acceptable practice. Of about 9 lbs. protein intake per day, an average cow will use 2 to 3 lbs. for milk protein synthesis. The difference of 1 lb. (which is the difference between a low- and a high-efficient cow), however, would make a huge difference on nitrogen losses and consequently ammonia emission from manure.



Protein (as amino acids) is an important nutrient for maintenance and production and its deficiency would lead to loss of production and poor animal health and reproduction. Protein requirements, however, may be over-predicted by the current dairy protein models, which leads to over-feeding of protein, deliberately through ration formulation, or due to inadequacy of monitoring of feeding management practices. *Excess die-*

tary protein results in increased nitrogen excretion with urine and consequently, ammonia volatilization losses from manure. Ammonia is an environmental and human health hazard, contributing to fine particulate matter formation, eutrophication of surface waters and nitrate contamination of ground waters, and soil acidity.

For this project, the Penn State Extension Dairy Team identified 12 small

(50 cows) or larger (550 cows), free-stall and tie-stall dairies in Central, Southeast, and Southwest PA with scrape, gravity-flow, or flush manure management systems. All dairies fed a total mixed ration (TMR). The team performed extensive forage and TMR sampling to verify crude protein (CP) levels at the beginning and throughout the project (Figure 1).

Project main objective:

Reduce whole-farm nitrogen inputs and ammonia emissions from Pennsylvania dairies while maintaining or improving profitability as measured through Income Over Feed Cost (IOFC)



Measuring barn floor ammonia emissions



PENNSTATE



Cooperative Extension
College of Agricultural Sciences

Fig. 1. Average TMR crude protein before (Fall'09 & Spring'10) and after (Fall'10 & Spring'11) protein reduction






Fig. 2. Barn floor ammonia emissions



temperature to more truly represent the effect of feeding, showed a **36% reduction in ammonia emission due to the reduced protein content of the TMR** (Figure 3). Overall, flush dairies emitted less ammonia off the barn floor than scrape or gravity-flow dairies.

Fig. 3. Laboratory ammonia emissions (Ammonia Emitting Potential of manure)



Following consultations with the dairymen and their nutritionists, a 1%-unit drop in TMR CP was achieved in most dairies in the 2nd year of the project (Figure 1). Barn floor and laboratory ammonia emissions were measured before and after the TMR protein reduction. So was milk yield per cow, milk composition (including milk urea nitrogen, MUN), and IOFC.

Following TMR protein reduction, barn floor ammonia emissions were reduced on average by about 65% (Figure 2), but a large portion of this difference was likely due to differences in ambient temperature, which was 41°F during the low-CP TMR periods vs. 57°F for the high-CP periods. Our laboratory measurements, conducted at the same temperature

to more truly represent the effect of feeding, showed a **36% reduction in ammonia emission due to the reduced protein content of the TMR** (Figure 3). Overall, flush dairies emitted less ammonia off the barn floor than scrape or gravity-flow dairies.

Contacts: In the Department of Dairy and Animal Science: Alex

Hristov, associate professor of dairy nutrition, 814-863-3669, anh13@psu.edu and Virginia Ishler, nutrient management specialist and Penn State Dairy Research Complex manager, 814-863-3912, vishler@psu.edu; and in Extension: Gene Schurman, dairy educator, 724-465-3880, exs10@psu.edu.

Laboratory measurement of the ammonia emitting potential of manure




Fig. 4. Income Over Feed Cost for Dairies Participating in the Project





Fig. 5. Average Milk Yield Before and After TMR Protein Content was Reduced



Conclusion: Reduction of dietary crude protein by about 1%-unit resulted in a significant 36% reduction in ammonia emissions from manure, did not affect milk yield, and increased income over feed costs by \$0.90/cow/d in commercial Pennsylvania dairies.

An **OUTREACH** program of the College of Agricultural Sciences

Penn State College of Agricultural Sciences research and extension programs are funded in part by Pennsylvania counties, the Commonwealth of Pennsylvania, and the U.S. Department of Agriculture.

Where trade names appear, no discrimination is intended, and no endorsement by Penn State Cooperative Extension is implied.

Penn State encourages persons with disabilities to participate in its programs and activities. If you anticipate needing any type of accommodation or have questions about the physical access provided, please contact 888-373-7232 in advance of your participation or visit.

This publication is available in alternative media on request.

The Pennsylvania State University is committed to the policy that all persons shall have equal access to programs, facilities, admission, and employment without regard to personal characteristics not related to ability, performance, or qualifications as determined by University policy or by state or federal authorities. It is the policy of the University to maintain an academic and work environment free of discrimination, including harassment. The Pennsylvania State University prohibits discrimination and harassment against any person because of age, ancestry, color, disability or handicap, national origin, race, religious creed, sex, sexual orientation, gender identity, or veteran status. Discrimination or harassment against faculty, staff, or students will not be tolerated at The Pennsylvania State University. Direct all inquiries regarding the nondiscrimination policy to the Affirmative Action Director, The Pennsylvania State University, 328 Boucke Building, University Park, PA 16802-5901; Tel 814-865-4700/V, 814-863-1150/TTY.