

### PLANT PATHOGENS

Disease Management

### Learning Objectives

- Become familiar with various types of plant pathogens or disease agents.
- Examine various methods of disease management.
- Familiarize student with common diseases for crop families in the midwest region

### Plant Pathogen Overview

- A plant pathogen is any harmful infectious agent, organism, or condition that reduces a plant's overall vitality, inhibits its growth, or limits the ability of the plant to survive and reproduce.
- Pathogens can be delivered in a multitude of different ways. These include: bacteria, fungi, viruses, nematodes, oomycetes, and abiotic toxicities.
- There are several different factors that need to be present in order to create an outbreak of disease on the farm. The pathogen must be present. There must be suitable host plants around. And there must be favorable environmental conditions for the growth and development of that particular plant pathogen.

- Bacteria
- Fungi
- Viruses
- Nematodes
- Abiotic
- Oomycetes

Bacteria: These single cell organisms absorb the nutrition from their host plant and thrive by killing the plant and living off of their decomposing organic matter. This is often described as a

saprophytic relationship. Bacterial pa

cause blights, rots, and wilts.

Onion rot caused by bacteria can be avoided by spacing bulbs closer. See

http://extension.umass.edu/vegetable/articles/stop-rotusing-cultural-practices-reduce-bacterial-bulb-decayonions

Fungi: Fungal pathogens are spread by many different environmental conditions such as wind, water, seeds, human and other non-human vectors. (A vector is a host capable of transferring a particular pathogen.) Fungi that are capable of regenerating spores during the growing season and re-infecting plants are known as polycyclic.

Fungi that must wait for next

Apple Scab is caused by a Fungus and can be treated with organic fungicides

https://www.extension.purdue.edu/extmedia/BP/BP-1-W.pdf



Viruses: Viruses are pieces of genetic material (RNA/DNA) that disturb the plants by mimicking naturally occurring genetic abnormalities. There are more than 700 plant viruses known. Viruses can be spread by tractors or equipment, tainted seed, or by a traveling vector.

Potato Virus Y is caused by a virus which is transmitted by aphids. For more information see <a href="http://labs.russell.wisc.edu/vegento/crops-and-insects/plant-pathogens/">http://labs.russell.wisc.edu/vegento/crops-and-insects/plant-pathogens/</a>

Nematodes: Nematodes are both a pathogen and a vector. Nematodes are microscopic worms and are one of the most abundant phyla with over 20,000 different species. A nematode either infects a plant by injecting a needle like mouth into the plant or it actually enters the plant with its entire body. The nematodes' saliva is the infecting agent and disturbs the metabolic process of the plant causing diseas

INFESTED

Carrots that have been diseased by nematodes. See

http://cetehama.ucanr.edu/?impact=718&a=7816

 Abiotic: This refers to deficiencies in the soil or surrounding environment that cause a debilitating

illness that is harmful or fatal



Iron deficiency in the soil. soil.



Magnesium deficiency in

Oomycetes: These organisms act much like a fungi, however they have a very different evolutionary history. They have mobile spores and can be primarily spread by both wind and water. They can also be spread by vectors. An example of an oomycetes is "downy mildew" also known as "damping off "

Oomycetes are responsible for this late blight. This is the same organism responsible for the Potato Famine and still has severe economic repercussions when it occurs today.



## Common Diseases in the Midwest See the videos detailing each

- blossom end rot <a href="http://youtu.be/6uObcfRWdK0">http://youtu.be/6uObcfRWdK0</a>
- early and late blight <a href="http://youtu.be/nqGODg8jhsl">http://youtu.be/9PabiuQ7wVI</a>
- powdery mildew <a href="http://youtu.be/9ZuTUiRD3Cs">http://youtu.be/9ZuTUiRD3Cs</a>
- downey mildew <a href="http://youtu.be/sz0vZ-t0gyg">http://youtu.be/sz0vZ-t0gyg</a>
- Scab <a href="http://youtu.be/tj8JCh4Why4">http://youtu.be/tj8JCh4Why4</a>
- peach curl <a href="http://youtu.be/9Wzu-aBVKdw">http://youtu.be/9Wzu-aBVKdw</a>

### Disease Management

 Plants often have developed natural defenses against pathogens. Some plants have developed disease tolerance. Some plants have developed disease resistance. Plants that are disease resistant will exhibit characteristics that actually repel certain harmful disease known to attack that particular plant. Plants that are disease tolerant can live with a problem pathogen but survive without any substantial reduction in yield or overall health.

## Disease Management Strategies

Proper greenhouse management can be extremely helpful in the control of outbreak and spread of pathogens. Disinfecting all soil trays and propagating mediums greatly reduces the incidence of pathogens.

See <a href="http://youtu.be/FzQyucZwj-E">http://youtu.be/FzQyucZwj-E</a>

### Disease Management Strategies

A solid approach to crop rotation will also limit the spread and overall effect of certain diseases. This is a result of diversity increasing overall vitality, limiting host species, and reducing environmental conditions needed for disease to flourish and spread in similar varieties or families of plants.

### Disease Management Strategies

Recognizing pre-existing factors that may contribute to pathogen outbreak is an important aspect of disease control. An example would be being vigilant and wary of downy mildew in a very wet or coastal environment.

Identifying the problem with the plant is the first step. See <a href="http://msue.anr.msu.edu/news/signs">http://msue.anr.msu.edu/news/signs</a> and symptoms of plant <a href="disease is it fungal viral or bacterial">disease is it fungal viral or bacterial</a> for information on distinguishing between viral, bacterial, and fungal diseases.

# Disease Management Strategies


In some organic situations, chemical control of disease is necessary and allowed by organic standards. Affected host plants can be physically removed or treated with such elements as copper, sulfur, or neem. It is always recommended that you consult with an expert before using or adding any amendment to your fields to help control disease.

### Disease Management Strategies

 Overall plant vitality is probably the single most important element for fighting and combating disease in your garden or field. Pathogens have the tendency to attack weak or stressed plants in the garden. If a disease establishes itself using the weaker plants it can often jump to the stronger crops. Therefore, maintaining conditions favorable to general plant vitality can significantly reduce the occurrence of disease. Removing harvested plants quickly and composting effectively will reduce instance of pathogens as well.

### OMRI Organic Materials Review Institute

Use OMRI's website to find appropriate products for disease management.



### Assessment/Review

- What is a vector?
- What are several types of pathogens and how do they affect the plant?
- What essential disease prevention methods should be used on the farm to reduce the risk of pathogens?

#### Resources

- https://ag.purdue.edu/btny/Pages/default.aspx
- http://msucares.com/insects/index.html
- http://msue.anr.msu.edu/news/signs\_and\_sym ptoms\_of\_plant\_disease\_is\_it\_fungal\_viral\_or bacterial