

Impacts of Patch-Burn Grazing and Winter-Patch Grazing on Arthropod Community Heterogeneity

Alyssa Vachino and Lan Xu*

Department of Natural Resource Management, South Dakota State University
Brookings, SD 57006



Introduction

- Historically, the Northern Great Plains evolved to thrive under periodic fire and substantial grazing pressures.
- Contemporary management practices focused on maximizing livestock production and fire suppression have resulted in more homogeneous plant communities.
- Arthropods are an important component of grassland systems and food webs.
- Arthropod community heterogeneity depends greatly on plant community heterogeneity.

Objectives

The objective of this study was to evaluate and compare the impact of patch-burn grazing and winter-patch grazing on arthropod community order richness, abundance, and diversity.

Hypothesis

We hypothesized that patch-burn grazing (PBG) and winter-patch grazing (WPG) will have different impacts on arthropod community compared to continuous season-long grazing (CG) in:

- Abundance
- Order Richness
- Diversity

Study Area

The location for this study was the Cottonwood Range and Livestock Research Station in western South Dakota (Lat. 43° 55'08" N, Long. 101° 52'58" W). The site is home to northern mixed-grass prairie vegetation and is mainly used for cattle grazing. Predominant soils are clayey and loamy. Wildfire burned part of each pasture in October 2016.




Fig. 1. Study design setup and execution

Methods

- Randomized Complete Block Design with 3 treatments (WPG, PBG, CG), 3 replicates (Pasture 3, 5, 6) for each treatment and 3 enclosures within each replicate (Fig. 1A)
- Four samples were taken from each enclosure in each pasture radiating out from the starting location of the enclosure in cardinal directions along 50-m transect (Fig. 1B)
- Samples were taken with a sweep net that was swung through the tops of the foliage in 180 degree swipes (Fig. 1C) and 70 swipes made along each transect (Fig. 1D)
- Each sample was transferred to a gallon labeled bag and stored in the freeze until sorting.
- Two sampling events occurred from early June to early August in 2018 two years post disturbance (PBG and WPG)
- Arthropods were sorted to taxonomic order.
- One-way ANOVA with significance $P < 0.05$

Results




Fig. 2. Total Arthropod abundance in CG, PBG, and WPG treatments




Fig. 4. Rank-abundance curves for Arthropod orders in CG, PBG, and WPG treatments




Fig. 3. Average Arthropod order richness in CG, PBG, and WPG treatments





Fig. 5. Arthropod community composition based on orders for CG, PBG, and WPG



Fig. 6. (A) Hemiptera, (B) Diptera, (C) Hymenoptera, (D) Lepidoptera, (E) Neuroptera, (F) Ephemeroptera, (G) Araneae, (H) Coleoptera, (I) Trichoptera, (J) Phasmoda, (K) Orthoptera, (L) Odonata

Results

- Sampling resulted in a total of 24,141 arthropods representing 12 orders (Fig. 6) across CG (5,122 arthropods, 9 orders), PBG (11,482 arthropods, 11 orders), and WPG (7,537 arthropods, 10 orders) treatments (Fig. 5).
- PBG and WPG significantly increased arthropods abundance compared to CG by 2.24 to 1.47 times, respectively (Fig. 2).
- No statistically significant difference was found in order richness among the 3 treatments, however, fewer orders were captured in CG treatments than in PBG or WPG treatments (Fig. 3).
- Three orders (Ephemeroptera, Neuroptera and Odonata) were only found in WPG and PBG treatments (Fig. 5), while the order Phasmoda was found only in CG treatment.
- Hemiptera, Diptera, and Hymenoptera were consistently the three most abundant orders in each treatment (Fig. 5) and made up 90% of abundance (Fig. 4)
- Rank-abundance curves appeared PBG and WPG have increased biodiversity compared to CG treatment (Fig. 4)



Discussion

- Results demonstrate that patch-burn grazing and winter-patch grazing has significantly positive impact on arthropods abundance.
- Patch-burn grazing and winter-patch grazing appears to harbor, on average, more orders (i.e. order richness).
- Patch-burn grazing may be preferable over CG for maintaining abundant and diverse Arthropod communities.
- WPG may serve as a better non-pyric management tool for enhancing ecosystem structure and services via food web diversity and pollination.

References

- Adler, P. D. Raff, W. & Lauenroth. 2001. The effect of grazing on the spatial heterogeneity of vegetation. *Oecologia* 128: 465-479
- BugGuide.net retrieved from <https://bugguide.net/node/view/15740>
- Collins, S. 1992. Fire frequency and community heterogeneity in tallgrass prairie vegetation. *Ecology* 73: 2001-2006
- Dennis, P. M. Young, & I. Gordon. 1998. Distribution and abundance of small insects and arachnids in relation to structural heterogeneity of grazed, indigenous grasslands. *Ecological Entomology*, 23: 253-264

Acknowledgements

This research was funded by the U.S. Department of Agriculture AFRI (Grant No. 2017-67020-26511) and NCR-SARE (Grant No. 2015-38640-23781), through a grant awarded to South Dakota State University, Joseph F. Nelson Undergraduate Research Mentorship Award, and the Griffith Undergraduate Research Award of South Dakota State University. Special thanks to Patrick Wagner, Madeline St. Claire, Luke Zilverberg, Jake Comer for their help in the lab and field.