

Fig. 3-2. Percent utilization in alternating rows of perennial ryegrass (PRG), meadow bromegrass (MB), orchardgrass (OG) and tall fescue (TF) with birdsfoot trefoil (BFT) in three stocking cycles. Overall mixed row utilization is shown for comparison. Shorter, darker bars show day 3 utilization, while taller, opaque bars show day 7 utilization. Graph D compares cycle 1 day 7 utilization (dark bars) with cycle 3 day 7 utilization (opaque bars). Bars within a grass-trefoil group followed by different letters are significantly different ($P=0.05$). Uppercase letters indicate differences on day 3 and lowercase letters indicate differences on day 7.




Fig. 3-3. Percent utilization in mixed rows of perennial ryegrass (PRG), meadow bromegrass (MB), orchardgrass (OG) and tall fescue (TF) with birdsfoot trefoil (BFT) in three stocking cycles. Shorter, darker bars show day 3 utilization, while taller, opaque bars show day 7 utilization). Graph D compares cycle 1 day 7 utilization (dark bars) with cycle 3 day 7 utilization (opaque bars). Bars within a grass-trefoil group followed by different letters are significantly different ($P=0.05$). Uppercase letters indicate differences on day 3 and lowercase letters indicate differences on day 7.