

Table 1. Effect of in-feed supplementation of CR and TC on relative liver weight of chickens fed with 2.5 µg/g AF. CR: carvacrol; TC: trans-cinnamaldehyde; AF: aflatoxins^{1,2}.

Items ³	Relative Liver Weight			
	Week 2	Week 3	Week 4	Week 5
CR				
Treatments				
Control	3.42% ± 0.71% ^a	2.50% ± 0.44% ^a	2.52% ± 0.34% ^a	2.44% ± 0.31% ^a
CR control	3.45% ± 0.90% ^a	2.83% ± 0.48% ^{ab}	2.51% ± 0.21% ^a	2.35% ± 0.27% ^a
AF	3.69% ± 0.41% ^a	3.04% ± 0.66% ^b	2.74% ± 0.30% ^a	2.73% ± 0.41% ^b
AF + CR	3.79% ± 0.34% ^a	2.65% ± 0.31% ^{ab}	2.59% ± 0.28% ^a	2.40% ± 0.19% ^a
TC				
Treatments				
Control	3.42% ± 0.71% ^a	2.50% ± 0.44% ^a	2.52% ± 0.34% ^{ab}	2.44% ± 0.31% ^a
TC control	3.49% ± 0.50% ^a	2.54% ± 0.32% ^a	2.53% ± 0.23% ^{ab}	2.41% ± 0.35% ^a
AF	3.69% ± 0.41% ^a	3.04% ± 0.66% ^b	2.74% ± 0.30% ^b	2.73% ± 0.41% ^b
AF + TC	3.53% ± 0.55% ^a	2.56% ± 0.51% ^a	2.35% ± 0.40% ^a	2.38% ± 0.09% ^a

^{a-b} Means with different superscripts in a column differ significantly ($P < 0.05$).

¹Means represent 5 birds per pen and two pens per treatment.

²Data are the mean ± SEM obtained from 5 birds per pen and two pens per treatment.

Error bar indicates SEM (n=10/treatment).

³ Treatments include Control: feed with no AF and no CR/TC supplementation; CR control: 0.75% carvacrol control; TC control: 0.75% trans-cinnamaldehyde control; AF: 2.5 µg/g aflatoxins; AF+CR: 2.5 µg/g aflatoxins + 0.75% carvacrol; AF+TC: 2.5 µg/g aflatoxins + 0.75% trans-cinnamaldehyde.