

Trial crop	Steep Time (days)	pH	N	P	K	Al	B	Ca	Cl	Co	Cu	Fe
<i>A. retroflexus</i>	3	5.3		1.40%	1.44%	1.91%	0.19%	0.03%		0%	0.00%	3.05%
<i>A. retroflexus</i>	5	5.2		2.24%	2.20%	2.69%	0.25%	0.02%		0%	0.00%	3.51%
<i>C. album</i>	3	5.2		1.61%	1.94%	6.39%	0.96%	0.01%		0%	1.65%	2.73%
<i>C. album</i>	5	5.3		2.13%	2.22%	6.91%	1.13%	0.01%		0%	0.74%	3.16%
<i>S. peregrinum</i>	3	5.8		1.43%	1.68%	0.00%	0.84%	0.26%		0%	0.56%	1.28%
<i>S. peregrinum</i>	5	4.9		1.43%	1.41%	0.00%	0.87%	0.27%		0%	0.31%	1.79%
<i>T. officinale</i>	3	4.9		1.78%	2.22%	0.00%	1.01%	1.20%		0%	0.37%	1.48%
<i>T. officinale</i>	5	4.8		1.57%	2.05%	1.11%	1.61%	1.42%		0%	0.26%	2.34%
<i>T. pratense</i>	3	4.8		0.90%	1.56%	0.69%	0.16%	0.73%		0%	0.30%	0.49%
<i>T. pratense</i>	5	4.9		1.11%	1.71%	0.48%	0.18%	0.76%		0%	0.30%	0.59%
<i>U. dioica</i>	3	8.2		2.18%	2.65%	1.23%	1.81%	1.76%		0%	6.72%	0.32%
<i>U. dioica</i>	5	4.9		3.35%	3.21%	1.36%	2.78%	2.04%		0%	1.62%	0.71%

Trial crop	Steep Time (days)	pH	I	Mg	Mn	Mo	Na	Ni	S	Se	Si	Zn
<i>A. retroflexus</i>	3	5.3		0.52%	0.31%	0%	2.47%		4.33%			0.45%
<i>A. retroflexus</i>	5	5.2		0.91%	0.51%	0%	4.31%		4.07%			0.70%
<i>C. album</i>	3	5.2		1.48%	0.96%	0%	3.90%		0.96%			1.11%
<i>C. album</i>	5	5.3		1.73%	1.10%	0%	12.50%		1.23%			1.46%
<i>S. peregrinum</i>	3	5.8		0.66%	0.56%	0%	1.16%		3.88%			0.73%
<i>S. peregrinum</i>	5	4.9		0.77%	0.61%	0%	1.09%		8.87%			0.69%
<i>T. officinale</i>	3	4.9		1.42%	1.16%	0%	0.93%		2.40%			0.48%
<i>T. officinale</i>	5	4.8		1.53%	1.44%	0%	1.31%		0.87%			0.63%
<i>T. pratense</i>	3	4.8		0.67%	0.85%	0%	6.34%		2.91%			0.69%
<i>T. pratense</i>	5	4.9		0.78%	0.96%	0%	7.99%		3.90%			0.89%
<i>U. dioica</i>	3	8.2		1.83%	2.21%	0%	9.32%		2.31%			3.13%
<i>U. dioica</i>	5	4.9		2.07%	2.98%	0%	4.64%		5.92%			1.25%

Table 11: Liquid nutrient analysis, expressed as percent of dried plant tissue nutrient concentrations. The highest percentage of nutrient carryover for each nutrient at 3 and 5 days is displayed in bold.