

**Table 4. 2016 Heirloom dry bean pre-harvest measurements, Alburgh, VT.**

Variety	Plant height	Lodged	Pod distance to ground	Pod disease
	cm	%	cm	%
Black Calypso	28.2	<b>0.00</b>	1.17	52.5
Hutterite Soup	28.5	7.50	1.17	22.5
Jacob's Cattle Gold	32.6*	<b>0.00</b>	0.75	30.0
Jacob's Cattle	32.4*	<b>0.00</b>	1.83	30.0
Kenearly Yellow Eye	34.9*	<b>0.00</b>	<b>2.33</b>	25.0
King of the Early	30.6	<b>0.00</b>	0.50	47.5
Lowe's Champion	36.3*	<b>0.00</b>	1.08	32.5
Lina Sisco	29.5	<b>0.00</b>	0.00	20.0
Light Red Kidney	<b>38.3*</b>	<b>0.00</b>	0.17	37.5
Marifax	32.2*	<b>0.00</b>	0.08	32.5
Orca	34.7*	<b>0.00</b>	0.25	22.5
Peregrin	34.2*	7.50	0.00	<b>7.50</b>
Raquel	31.3	<b>0.00</b>	0.08	32.5
Spanish Tolasna	34.0*	<b>0.00</b>	0.83	37.5
Tongues of Fire	25.0	<b>0.00</b>	0.00	40.0
Tiger's Eye	31.5	<b>0.00</b>	0.00	30.0
Vermont Appaloosa	33.3*	<b>0.00</b>	0.00	35.0
Vermont Cranberry	27.4	<b>0.00</b>	0.00	42.5
<i>LSD (0.10)</i>	6.15	NS	NS	NS
<i>Trial Mean</i>	31.9	0.83	0.57	32.1

Values shown in **bold** are of the highest value or top performing.

\* Dry beans that did not perform significantly lower than the top performing variety in a particular column are indicated with an asterisk.

NS-Treatments were not significantly different from one another.