

Trial Crop	N	P	K	Al	B	Ca	Cl	Co	Cu	Fe
<i>Amaranthus retroflexus</i>		3.9	46.1	0.0022	0.096	6.9		0	0.6	0.0018
<i>Chenopodium album</i>		3.7	47.8	0.0007	0.0735	4		0	0.6	0.0014
<i>Sympytum peregrinum</i>		3.7	68.1	0.0013	0.1315	6.2		0	1.1	0.0022
<i>Taraxacum officinale</i>		5	36.1	0.0007	0.0654	2.7		0	1.2	0.0021
<i>Trifolium pratense</i>		2.6	19.1	0.0011	0.0597	2.8		0	0.5	0.0026
<i>Urtica dioica</i>		4	24.4	0.0005	0.0808	8.1		0	0.6	0.0018

Trial Crop	I	Mg	Mn	Mo	Na	Ni	S	Se	Si	Zn
<i>Amaranthus retroflexus</i>		1.9	0.2269	1.5	2.2	0	7.4			0.7303
<i>Chenopodium album</i>		1.3	0.1243	1.6	0.5	0	6			0.5987
<i>Sympytum peregrinum</i>		1.2	0.1145	1	5.6	0	3.8			0.3576
<i>Taraxacum officinale</i>		0.8	0.0612	2	8.6	0	7.2			0.5776
<i>Trifolium pratense</i>		0.8	0.0398	1.8	1.2	0	3.3			0.2925
<i>Urtica dioica</i>		1.3	0.1266	2.1	0.8	0	6			0.2295

Table 8: Bioaccumulation factor averages for 6 trial crops.